




Phyt. 926 $\frac{L}{1}$

15 Bde in 17 Kplt.

Mit den Erg. Auctarium, Floral Reg.

(Temmer No. 159 Bde. 1-5 850.-)

(4 No. 257 Auk. 35 1863 1-6 1150.-! 1500)

1. Ausgabe. 1825 f.

Nissen 2222, Pritzel 6004

NEI 94 — *Maund, B.* The Botanic Garden, consisting of highly finished Representations of hardy ornamental flowering Plants. Band 1—13. Floral Register 2 Bde. The Auctarium 2 Bde. The Fruitist. 3 Bde. London 1825 bis 1850. 8°. Mit 1248 kolorierten Pflanzenkupfern auf 312 Tafeln, 72 kol. Obstkupfern. Bd. 1—13 in grünen Halblederbänden d. Zt. mit roten. Rsch. und Blumenstempeln auf Buchrücken. Erg. Bde. in grün. Halbmaroquinbdn. d. Zt. Teilw. leicht berieben. 1800,—

Nissen 2222. Pritzel 6004. Durchweg sauberes Exemplar in hübschen Einbänden. In dieser Vollständigkeit sehr selten!

34-888

Qu. 9.60
LXX

14 vols (including *Summit*)

in 13 vols

+ Floral Register
x Auctioneer

Komplett aller Erscheinungen

BOTANIC GARDEN

HERBARIUM

GEORGE ENGELMANN PAPERS

1844

1845

1846

1847

1848

1849

1850

1851

1852

1853

C

THE
BOTANIC GARDEN;

CONSISTING OF
HIGHLY FINISHED REPRESENTATIONS
OF HARDY
ORNAMENTAL FLOWERING PLANTS,
CULTIVATED
IN GREAT BRITAIN;

WITH
THEIR NAMES, CLASSES, ORDERS, HISTORY, QUALITIES, CULTURE,
AND PHYSIOLOGICAL OBSERVATIONS.

BY
Benjamin
B. MAUND, F.L.S.

PART I.

“Not a tree,
A plant, a leaf, a blossom, but contains
A folio volume. We may read and read,
And read again, and still find something new,
Something to please, and something to instruct.”
HURDIS.

London:

PUBLISHED BY SIMPKIN & MARSHALL, STATIONERS' HALL COURT.

1825.

Wb/64/674

THE
BOTANIC GARDEN

CONSISTING OF

NIGHTLY VIGNETTE REPRESENTATIONS

OF PLANTS

ORIGINATING IN THE PLANTS

OF GREAT BRITAIN

IN GREAT BRITAIN

Bayerische
Staatsbibliothek
München

R. MAUND, F.R.S.


PART I.

"Not a tree,
A plant, a leaf, a flower, but a tree,
A little tree, we say, and yet
And yet again, and still the same,
Resisting to the wind and weather's power."
Herrick.

LONDON:

PRINTED BY WILKINSON & MARRIAGE, STATIONERS, BELL COURT.

1865.



To
Her Most Gracious Majesty.
The Queen.

In grateful acknowledgment
of her
Condescension and Patronage.
this Volume

Is most respectfully
Dedicated.

By
Her Majesty's
Most loyal, obliged,
And devoted, Servant.

Benjamin Mownd.

TO
THE LOVERS OF BOTANY,
IN PARTICULAR,
AND
THE ADMIRERS OF A FLOWER GARDEN,
GENERALLY,
THIS ATTEMPT
TO INCREASE THEIR GRATIFICATIONS,
IS
MOST RESPECTFULLY INSCRIBED.

THE LOVERS OF BOTANY

IN PROSE

AND

THE ADORNERS OF A FLOWER GARDEN

BY

THIS ATTEND

TO EXHIBIT THEIR ORIGINATIONS

IN

MOST RESPECTFULLY INSCRIBED

PREFACE.

Man, by nature, inherits the love of flowers. The domains of the noble, and the cottage of the humble, alike proclaim to us the dominion of this passion.

If the busy scenes of life, in which many are obliged to move, suppress for awhile this divine excitement, yet the chief occupant of their thoughts is no sooner relinquished, than natural inclinations immediately evince themselves. They sigh for rural retirement, there to enjoy the uncontaminated atmosphere of nature, to cultivate its choicest gifts, and to linger over its vegetable beauties.

The attraction of a flower garden, the health yielded by its cultivation, and the ten thousand gratifications arising out of it, are freely acknowledged by every one. Open as these manifold pleasures are, to the least individual amongst us, the author of the Botanic Garden was desirous of producing a work, at such moderate price, as none may be excluded from possessing. One that should disseminate information amongst the lovers of flowers, encourage the taste for their cultivation, and yield a stimulus to the exertions of those who duly appreciate the enjoyments that invariably arise from so pure a source.

Whether he has effected the desired object, is left to the decision of his readers. The best exertions of his humble ability have been employed; and the encouragement of the public has laid him under obligations to continue those efforts with redoubled zeal.

Much of a work like the present must necessarily be dependant on the labours of preceding botanists; on men who have toiled, and some who still toil, to raise to perfection a delightful science, inexhaustible in extent and intricate in its ramifications. In this little work it is not the province of the author to wade deep in the current of science and research. He humbly collects from the sweets and the beauties that float on the surface, and of these he has pleasure in composing a nosegay, as a periodical present to the lovers of a flower garden.

Many ideas, it is presumed, will be found in the following sheets, which have arisen from the author's own experience. Others may be met with, which for the hundredth time are presented to the eye of the reader. But whether old or new, original or selected, his endeavour will be to keep in view one principal object,—the production of every useful and interesting information connected with the subjects on which he treats.

Regarding the correctness of the plates he can speak with confidence. Talented artists have lent their aid, and every exertion has been made to render them as perfect portraits as the state of the arts, and pecuniary remuneration, will admit.

That he owes much gratitude to several individuals of noble rank, for their condescension in forwarding his views, he is deeply sensible. He would have pleasure in a more explicit acknowledgment of their favours; but that superior minds feel no gratification in whatever may assume the form of adulation.

The reception of the Botanic Garden can but inspire the author with additional zeal in his favourite pursuit, and stimulate his endeavours to secure a continuance of that patronage, which in the present excess of literary productions, may be deemed no trifling distinction.

INDEX TO VOLS. I to VI.

<i>Systematic Name.</i>	<i>English Name.</i>	<i>Vol.</i>	<i>No.</i>
<i>Acanthus spinosus</i> ,	Prickly-leaved Acanthus, ..	6	508
<i>Achillea clavennæ</i> ,	Silver-leaved Milfoil,	2	122
<i>Achillea speciosa</i> ,	Spear-leaved Milfoil,	1	44
<i>Aconitum napellus</i> ,	Monk's Hood,	3	210
<i>Aconitum versicolor</i> ,	Shaded Monk's-Hood,	5	436
<i>Adonis vernalis</i> ,	Spring Adonis,	1	27
<i>Æthionema membranaceum</i> ,	Memb-podded Æthionema, ..	4	334
<i>Agrostemma suecica</i> ,	Swedish Agrostemma,	6	576
<i>Alstrœmeria pelegrina</i> ,	Spotted Alstrœmeria,	5	435
<i>Alyssum saxatile</i> ,	Rock Madwort,	3	254
<i>Amaryllis lutea</i> ,	Yellow Amaryllis,	1	2
<i>Ammobium alatum</i> ,	Wing-stalked Ammobium ..	3	219
<i>Amsonia latifolia</i> ,	Broad-leaved Amsonia,	1	18
<i>Anagallis fruticosa</i> ,	Shrubby Anagallis,	6	485
<i>Anagallis Monelli</i> ,	Monelli's Pimpernel,	5	399
<i>Anagallis Webbiana</i> ,	Webb's Pimpernel,	4	343
<i>Anchusa Italica</i> ,	Italian Buglos,	2	173
<i>Andromeda calyculata</i> ,	Calycled Andromeda,	1	20
<i>Andromeda coriacea</i>	Leather-leaved Andromeda, ..	2	171
<i>Andromeda Mariana</i> ,	Maryland Andromeda,	3	240
<i>Anemone coronaria</i> ,	Poppy Anemone,	2	103
<i>Anemone hortensis</i> ,	Garden Anemone,	2	191
<i>Anemone nemorosa</i> , flôr. plen.	Double Wood Anemone,	4	318
<i>Anemone palmata</i> ,	Palmated Anemone,	2	145
<i>Anemone pulsatilla</i> ,	Pasque Flower,	3	198
<i>Antennaria dioica</i> ,	Diœcious Antennaria,	3	247
<i>Antennaria triplinervis</i> ,	Three-nerved Antennaria, ..	6	504
<i>Anthericum liliastrum</i> ,	Savoy Spider-wort,	1	88
<i>Antirrhinum majus</i> ,	Great Snap-Dragon,	3	279
<i>Apocynum androsæmifolium</i> , ..	Tutsan-leaved Dog's-bane, ..	2	99
<i>Aponogeton distachyon</i> ,	Two-spiked Aponogeton, ..	6	559
<i>Aquilegia Canadensis</i> ,	Canadian Columbine,	4	322
<i>Arbutus unedo</i> ,	Common Strawberry Tree, ..	4	361
<i>Argemone Mexicana</i> ,	Mexican Argemone,	1	85
<i>Arnica montana</i> ,	Mountain Arnica,	3	239
<i>Asclepius tuberosa</i> ,	Tuberous Swallow-wort,	1	78
<i>Asphodelus luteus</i> ,	Yellow King's-spear,	3	246
<i>Asphodelus ramosus</i> ,	Branched King's-spear,	3	251
<i>Aster alpinus</i> ,	Alpine Aster,	2	124
<i>Aster amellus</i> ,	Italian Starwort,	2	188
<i>Aster Novæ-Anglæ, ruber</i> , ..	Red New England Starwort, ..	3	223
<i>Aster Sibiricus</i> ,	Siberian Aster,	5	411
<i>Aster tenellus</i> ,	Slender Aster,	3	238
<i>Astragalus alopecuroides</i> , ..	Fox-tail-like Milk Vetch, ..	3	265
<i>Astragalus monspessulanus</i> , ..	Montpelier Milk Vetch,	6	531
<i>Astrantia major</i> ,	Greater Black Masterwort, ..	3	212
<i>Astrantia maxima</i> ,	Three-leav. Bk. Masterwort, ..	3	213
<i>Atropa belladonna</i> ,	Deadly Nightshade,	2	105

<i>Systematic Name.</i>	<i>English Name.</i>	<i>Vol.</i>	<i>No.</i>
Aubrietia purpurea,	Tufted Aubrietia,	3	230
Azalea calendulacea,	Marygold-like Azalea,	4	313
Azalea nudiflora,	Naked-flowered Azalea,	6	541
Azalea pontica,	Yellow, or Pontic Azalea, ..	3	261
Baptisia Australis.	Southern Baptisia,	6	567
Berberis vulgaris,	Common Berberry,	4	383
Betonica incana,	Hoary Betony,	4	312
Bignonia radicans,	Ash-leaved Trumpet-flower, 2		123
Biscutella hispida,	Hispid Biscutella,	3	206
Bouvardia triphylla,	Three-leaved Bouvardia, ..	4	384
Briza maxima,	Large Quaking-grass,	4	382
Buddlea globosa,	Round-headed Buddlea,	1	58
Bulbocodium-vernum,	Spring Bulbocodium,	6	557
Bupthalmum grandiflorum, ..	Great-flowered Ox-eye,	2	155
Cacalia coccinea,	Scarlet-flowered Cacalia, ..	1	76
Calampelis scabra,	Rough-podded Calampelis, ..	4	289
Calceolaria arachnoidea, ..	Cobweb Slipperwort,	4	357
Calceolaria corymbosa,	Chili Slipperwort,	1	67
Calceolaria pinnata,	Wing-leaved Slipperwort, ..	2	166
Calceolaria plantaginea,	Plantain-leaved Slipperwort, 4		328
Calendula pluvialis,	Small Cape Marygold,	3	280
Calliopsis Tinctoria,	Arkansa Calliopsis,	6	538
Calluna vulgaris, flore pleno.	Double-flowered Ling,	3	266
Campanula Carpatica,	Carpathian Bell-flower,	2	130
Campanula excisa,	Cut Bell-flower,	6	490
Campanula glomerata,	Clustered Bell-flower,	2	144
Campanula grandiflora,	Great-flowered Bell-flower, 6		516
Campanula hederacea,	Ivy-leaved Bell-flower,	6	496
Campanula lactiflora,	Milk-coloured Bell-flower, ..	3	200
Campanula nitida,	Smooth-leaved Bell-flower, 3		224
Campanula persicifolia,	Peach-leaved Bell-flower, ..	2	164
Campanula pumila,	Dwarf Bell-flower,	1	48
Campanula pyramidalis,	Pyramidal Bell-flower,	4	315
Campanula speciosa,	Showy Bell-flower,	3	228
Campanula speculum,	Venus's Looking-glass,	2	147
Caprifolium pubescens,	Pubescent Honeysuckle,	6	565
Carthamus tinctorius,	Officinal Carthamus,	2	162
Catananche cœrulea,	Blue flowered Catananche, ..	1	15
Catananche cœrulea,	Two-coloured Catananche, ..	5	424
Ceanothus intermedia,	Intermediate Red-Wood, ..	6	572
Celsia Critica,	Cretan Celsia,	5	400
Centaurea suaveolens,	Yellow Sultan,	1	95
Centaurea glastifolia,	Woad-leaved Centaury,	6	506
Centaurea ochroleuca,	Ivory-coloured Centaury, ..	5	462
Centaurea crocodylium,	Blush Centaury,	5	431
Centrocarrha chrysomela, ..	Yel. and Dark Centrocarrha,	6	528
Cerastium Biebersteinii,	Bieberstein's Cerastium,	6	515
Cheiranthus Alpinus,	Alpine Wall Flower,	6	518

<i>Systematic Name.</i>	<i>English Name.</i>	<i>Vol.</i>	<i>No.</i>
Cheiranthus tenuifolius,	Fine-leaved Wall-flower, ..	6	558
Chelone barbata,	Scarlet Chelone,	1	81
Chelone obliqua,	Red-flowered Chelone,	1	92
Chrysanthemum coronarium, ..	Garden Chrysanthemum, ..	2	143
Chrysanthemum Sinense, ..	Chinese Chrysanthemum, ..	2	120
Chrysanthemum Sinense, ..	Chinese Chrysanthemum, ..	4	353
Chrysanthemum tricolor, ..	Three-col. Chrysanthemum, ..	1	91
Chrysocoma linosyris,	Flax-leaved Goldyllocks, ..	4	358
Chymocarpus pentaphyllus, ..	Five-leaved Chymocarpus, ..	5	480
Cineraria macrophylla,	Sea Ragwort,	5	451
Cineraria maritima,	Large-leaved Cineraria,	6	524
Circæa alpina,	Alpine Nightshade,	4	378
Cistus acutifolius,	Acute leaved Rock Rose, ..	5	449
Clarkia pulchella,	Fair-flowering Clarkia,	3	199
Claytonia Sibirica,	Siberian Claytonia,	4	344
Claytonia Virginica,	Virginian Claytonia,	2	138
Clematis integrifolia,	Entire-leav. Virgin's-bower, ..	2	98
Cochlearia Grænländica, ..	Greenland Scurvy-grass, ..	1	64
Collinsia grandiflora,	Large-flowered Collinsia, ..	5	391
Collomia grandiflora,	Large-flowered Collomia, ..	5	392
Collomia linearis,	Linear-leaved Collomia,	5	417
Colutea frutescens,	Scarlet Bladder Senna,	1	37
Commelina tuberosa,	Tuberous-root. Commelina, ..	3	257
Coreopsis Atkinsoniana,	Atkins's Coreopsis,	5	389
Coreopsis lanceolata,	Spear-leaved Coreopsis,	1	46
Coreopsis tenuifolia,	Slender-leaved Coreopsis, ..	1	62
Coreopsis tinctoria,	Arkansa Coreopsis,	1	13
Coreopsis tripteris,	Three-leaved Coreopsis,	4	352
Coreopsis verticillata,	Whorl leaved Coreopsis,	6	562
Cornus Canadensis,	Canadian Dogwood,	2	136
Coronilla Iberica,	Iberian Coronilla,	5	446
Coronilla varia,	Various-coloured Coronilla, ..	3	226
Cortusa Mathioli,	Mathioli's Sanicle,	4	296
Cosmea bipinnata,	Double-pinnate-leav. Cosmea, ..	6	534
Crocus Susianus,	Cloth of Gold Crocus,	2	121
Crocus versicolor,	Party-coloured Crocus,	2	151
Cryptostemma calendulaceum, ..	Marygold-flo. Cryptostemma, ..	5	459
Cuscuta verrucosa,	Wart-calyxed Dodder,	1	10
Cyclamen coum,	Round-leaved Cyclamen, ..	3	229
Cynoglossum omphalodes, ..	Comfrey-lvd Hound's-tongue, ..	3	243
Cytisus argenteus,	Silver-leaved Cytisus,	5	452
Cytisus capitatus,	Headed Cytisus,	1	90
Cytisus nigricans,	Black-rooted Cytisus,	2	149
Dahlia superflua,	Dahlia, Blood-red,	4	297
Dahlia superflua,	Dahlia, Crimson Globe,	4	306
Dahlia superflua,	Dahlia, Painted Lady,	4	329
Dahlia superflua,	Double Purple Dahlia,	2	115
Daphne cneorum,	Trailing Daphne,	3	256

<i>Systematic Name.</i>	<i>English Name.</i>	<i>Vol.</i>	<i>No.</i>
<i>Daphne gnidium</i> ,	Flax-leaved Daphne,	1	72
<i>Daphne tarton-raira</i> ,	Silvery-leaved Daphne,	2	110
<i>Delphinium consolida</i> ,	Branching Larkspur,	5	432
<i>Delphinium grandiflorum</i> , ..	Large-flowered Larkspur, ..	1	29
<i>Delphinium mesoleucum</i> , ..	White-centred Larkspur, ..	5	403
<i>Dianthus caryophyllus</i> ,	Carnation,	4	301
<i>Dianthus caryophyllus</i> ,	Prince of Orange Picotee, ..	3	287
<i>Dianthus caryophyllus</i> ,	Yellow Picotee,	2	137
<i>Dianthus Chinensis</i> ,	China Pink,	1	65
<i>Dianthus deltoides</i> ,	Maiden Pink,	2	142
<i>Dianthus Hispanicus</i> ,	Spanish Pink,	2	108
<i>Dianthus Japonicus</i> ,	Japanese Pink,	2	168
<i>Dianthus plumarius</i> ,	Feathered Pink,	3	253
<i>Dianthus superbus</i> ,	Superb Pink,	5	425
<i>Diclytra eximia</i> ,	Choice Diclytra,	4	340
<i>Dictamnus fraxinella</i> ,	Red Fraxinella,	4	307
<i>Digitalis hybrida</i> ,	Campbell's Hybrid Foxglove, ..	5	468
<i>Digitalis laciniata</i> ,	Cut-leaved Foxglove,	5	447
<i>Digitalis lutea</i> ,	Small Yellow Foxglove,	1	94
<i>Digitalis minor</i> ,	Dwarf Foxglove,	4	299
<i>Diplocoma villosa</i> ,	Villous Diplocoma,	6	529
<i>Dodecatheon Meadia</i> ,	American Cowslip,	1	25
<i>Dolichos lignosus</i> ,	Woody Dolichos,	5	405
<i>Draba hirta</i> ,	Hairy Whitlow-grass,	2	132
<i>Dracocephalum Altaianse</i> , ..	Altaiian Dragon's Head,	6	563
<i>Dracocephalum canescens</i> , ..	Canescent Dragon's Head, ..	6	502
<i>Dracocephalum Moldavicum</i> , ..	Moldavian Dragon's Head, ..	4	371
<i>Dracocephalum speciosum</i> , ..	Showy Dragon's Head,	1	57
<i>Dryas octopetala</i> ,	Eight-petaled Dryas,	6	566
<i>Epilobium Dodonæi</i> ,	Dodonæus's Epilobium,	1	55
<i>Erica australis</i> ,	Spanish Heath,	1	54
<i>Erica ciliaris</i> ,	Ciliated Heath,	6	494
<i>Erica cinerea</i> ,	Grey Heath,	5	434
<i>Erica herbacea</i> ,	Early Dwarf Heath,	1	22
<i>Erica mediterranea</i> ,	Mediterranean Heath,	1	74
<i>Erica multiflora</i> ,	Many-flowered Heath,	4	376
<i>Erica stricta</i> ,	Straight-branched Heath, ..	3	217
<i>Erigeron asteroides</i> ,	Aster-like Erigeron,	5	453
<i>Erinus alpinus</i> ,	Alpine Erinus,	1	11
<i>Erodium hymenodes</i> ,	Ternate-leaved Heron's-Bill, ..	3	193
<i>Erpetion reniformis</i> ,	Spurless Violet,	4	342
<i>Eryngium alpinum</i> ,	Alpine Eryngo,	5	428
<i>Eryngium Bourgati</i> ,	Bourgati's Eryngo,	5	396
<i>Erythræa aggregati</i> ,	Aggregate Erythræa,	4	341
<i>Erythronium Americanum</i> , ..	Yel-flow. Dog's-tooth Violet, ..	2	178
<i>Erythronium dens canis</i> ,	Common Dog's-tooth Violet, ..	2	181
<i>Escallonia rubra</i> ,	Red-flowered Escallonia, ..	6	553
<i>Eupatorium maculatum</i> ,	Spotted-stalked Eupatorium, ..	2	107

<i>Systematic Name.</i>	<i>English Name.</i>	<i>Vol.</i>	<i>No.</i>
<i>Eschscholtzia Californica</i> , ..	Californian <i>Eschscholtzia</i> , ..	4	298
<i>Entoca viscida</i> ,	Clammy <i>Eutoca</i> ,	6	569
<i>Fragaria Indica</i> ,	Yel.-flowered Strawberry, ..	1	7
<i>Fritillaria meleagris</i> ,	Chequered <i>Fritillaria</i> ,	2	183
<i>Fuchsia macrostema</i> ,	Globe-flowered <i>Fuchsia</i> ,	6	533
<i>Fumaria Halleri</i> ,	Haller's <i>Fumatory</i> ,	3	262
<i>Fumaria lutea</i> ,	Yellow-flowered <i>Fumatory</i> , ..	2	111
<i>Fumaria nobilis</i> ,	Great-flowered <i>Fumatory</i> , ..	1	69
<i>Gailardia aristata</i> ,	Awed <i>Gailardia</i> ,	5	402
<i>Gailardia bicolor</i> ,	Two-coloured <i>Gailardia</i> , ..	2	100
<i>Galega Persica</i> ,	Persian Goat's Rue,	5	455
<i>Galinsogea trilobata</i> ,	Three-lobed <i>Galinsogea</i> , ..	4	370
<i>Gaultheria procumbens</i> ,	Trailing <i>Gaultheria</i> ,	1	17
<i>Gaura biennis</i> ,	Biennial <i>Gaura</i> ,	1	75
<i>Gazania rigens</i> ,	Rigid <i>Gazania</i> ,	5	415
<i>Genista ephedroides</i> ,	Ephedra-like <i>Genista</i> ,	6	498
<i>Genista sagittalis</i> ,	Jointed <i>Genista</i> ,	1	50
<i>Gentiana acaulis</i> ,	<i>Gentianella</i> ,	1	51
<i>Gentiana asclepiadea</i> ,	Swallow-wort-leav. <i>Gentian</i> , ..	3	282
<i>Gentiana Catesbæi</i> ,	Catesby's <i>Gentian</i> ,	2	113
<i>Gentiana cruciata</i> ,	Crosswort, or crossed <i>Gentian</i> , ..	5	387
<i>Gentiana verna</i> ,	Spring <i>Gentian</i> ,	2	179
<i>Geranium anemonefolium</i> , ..	Anemone-leaved Crane's Bill, ..	6	545
<i>Geranium angulatum</i> ,	Angular-stalked Crane's Bill, ..	6	484
<i>Geranium Ibericum</i> ,	Iberian Crane's Bill,	5	458
<i>Geranium Lancastriense</i> ,	Lancaster Crane's Bill,	5	388
<i>Geranium pratense</i> , var.	Varieg. Meadow Crane's Bill, ..	6	492
<i>Geum Chilense</i> ,	Chile <i>Geum</i> ,	3	273
<i>Gilia capitata</i> ,	Cluster-flowered <i>Gilia</i> ,	3	202
<i>Gilia tricolor</i> ,	Three-coloured <i>Gilia</i> ,	6	509
<i>Gladiolus Natalensis</i> ,	Natal Corn-flag,	6	536
<i>Glaucium phœniceum</i> ,	Red-flowered Horn-Poppy, ..	4	349
<i>Globularia vulgaris</i> ,	Common <i>Globularia</i> ,	1	9
<i>Gnaphalium arenarium</i> ,	Sand Everlasting,	1	38
<i>Gratiola aurea</i> ,	Golden Hedge-Hyssop,	4	374
<i>Gypsophila prostrata</i> ,	Trailing <i>Gypsophila</i> ,	2	150
<i>Hedysarum obscurum</i> ,	Creeping <i>Hedysarum</i> ,	5	441
<i>Hedysarum roseum</i> ,	Rose-coloured <i>Hedysarum</i> , ..	4	346
<i>Helianthemum Marifolium</i> , ..	Marum-leaved Sun Rose, ..	6	574
<i>Helianthemum roseum</i> ,	Rose-coloured Sun Rose, ..	3	267
<i>Heliophila araboides</i> ,	Arabis-like <i>Heliophila</i> ,	6	522
<i>Helleborus niger</i> ,	Christmas Rose,	1	8
<i>Helonius bullata</i> ,	Spear-leaved <i>Helonius</i> ,	3	235
<i>Hepatica trilobata</i> ,	Three-lobed <i>Hepatica</i> ,	4	323
<i>Hesperis matronalis</i> , alba, ..	Double White Rocket,	1	39
<i>Hesperis matronalis</i> , pur.	Double Purple Rocket,	1	70
<i>Hibiscus Syriacus</i> ,	<i>Althæa Frutex</i> ,	1	77
<i>Horminum Pyrenaicum</i> ,	Pyrenean <i>Horminum</i> ,	4	320

<i>Systematic Name.</i>	<i>English Name.</i>	<i>Vol.</i>	<i>No.</i>
Houstonia cærulea,	Blue-flowered Houstonia, ..	6	493
Hunnemannia fumariæfolia, ..	Fumitory-leav. Hunnemannia, ..	5	423
Hyacinthus orientalis,	Oriental Hyacinth,	4	367
Hyoscyamus niger,	Common Henbane,	2	172
Hypericum Kalmianum,	Kalmia-leav. St. John's wt. ..	3	194
Hypoxis erecta,	Upright Hypoxis,	4	302
Hyssopus Orientalis,	Oriental Hyssop,	5	466
Iberis sempervirens,	Narrow-leaved Candy-tuft, ..	1	82
Iberis Tenoreana,	Tenore's Candy-tuft,	6	556
Iris neglecta,	Horneman's Iris,	6	519
Iris pallida,	Pale Iris,	4	303
Iris pumila,	Dwarf Iris,	3	263
Iris Ruthenica,	Russian Iris,	5	456
Iris sambucina,	Elder-scented Iris,	5	443
Iris Sibirica,	Siberian Iris,	3	274
Iris Susiana,	Chalcedonian Iris,	1	30
Iris variegata,	Variegated Iris,	3	278
Iris versicolor,	Changeable Iris,	1	3
Iris Virginica,	Virginian Iris,	4	327
Isotoma axillaris,	Axillary-flowered Isotoma, ..	3	218
Jasione perennis,	Perennial Sheep's Scabious, ..	5	471
Jasminum revolutum,	Curled-flowered Jasmine, ..	1	12
Kalmia glauca,	Glaucous Kalmia,	1	43
Kalmia latifolia,	Broad-leaved Kalmia,	1	33
Kaulfussia amelloides,	Amellus-like Kaulfussia, ..	2	169
Lablab vulgaris,	Common Lablab,	5	404
Lachenalia tricolor,	Three-coloured Lachenalia, ..	1	21
Lathyrus Armitageanus,	Armitage's Lathyrus,	6	526
Lathyrus odoratus,	Sweet Pea,	4	369
Lathyrus rotundifolius,	Round-leaved Lathyrus,	6	511
Lathyrus tuberosus,	Tuberous Lathyrus,	2	139
Lavatera salvitellensis,	Pyramidal Lavatera,	6	505
Lavatera triloba,	Three-lobed Lavatera,	5	437
Ledum buxifolium,	Box-leaved Ledum,	1	52
Ledum latifolium,	Broad-leaved Ledum,	3	227
Ledum palustre,	Marsh Ledum,	1	47
Leptostelma maxima,	Great Leptostelma,	5	467
Leucocarpus alatus,	Wing-stalked Leucocarpus, ..	4	355
Leucojum æstivum,	Summer Snowflake,	2	186
Liatris scariosa,	Scaly-cupped Liatris,	2	161
Libertia formosa,	Beautiful Libertia,	5	472
Lilium bulbiferum,	Bulb-bearing Lily,	1	31
Lilium Canadense,	Canadian Lily,	2	112
Lilium Chalcedonicum,	Scarlet Martagon,	4	292
Lilium longiflorum,	Long-flowered Lily,	6	532
Lilium martagon,	Purple Martagon,	4	332
Lilium martagon,	White Martagon,	5	426
Lilium Pomponium,	Pompone Lily,	5	444

<i>Systematic Name.</i>	<i>English Name.</i>	<i>Vol.</i>	<i>No.</i>
<i>Lilium tigrinum</i> ,.....	Tiger-spotted Lily,.....	1	53
<i>Linaria alpina</i> ,	Alpine Toad-flax,	4	305
<i>Linaria bipartita</i> ,	Two-parted Toad-flax,	3	272
<i>Linaria cymbalaria</i> ,.....	Ivy-leaved Toad-flax,	4	300
<i>Linaria Dalmatica</i> ,	Dalmatian Toad-flax,	6	570
<i>Linaria Italica</i> ,	Italian Toad-flax,	5	463
<i>Linaria purpurea</i> ,	Purple Flax-weed,.....	1	34
<i>Linaria triornithophora</i> ,	Three-birds-bearing Td. Flax	5	440
<i>Linum alpinum</i> ,	Alpine Flax,	1	14
<i>Linum flavum</i> ,.....	Yellow Flax,	6	539
<i>Linum monogynum</i> ,	One-styled Flax,.....	6	482
<i>Lithospermum purpu. cerul.</i> , ..	Purple and blue Gromwell, ..	5	413
<i>Loasa acanthifolia</i> ,	Acanthus-leaved Loasa,.....	3	252
<i>Lobelia bicolor</i> ,	Two-coloured Lobelia,	4	351
<i>Lobelia fulgens</i> ,	Fulgent Lobelia,.....	1	73
<i>Lobelia fulgens-cærulea</i> , ..	Hybrid Lobelia,	6	535
<i>Lobelia syphilitica</i> ,.....	Blue Lobelia,	1	61
<i>Lobelia tupa</i> ,	Tupa Lobelia,	6	503
<i>Lonicera Tartarica</i> ,.....	Tartarian Honeysuckle,	1	23
<i>Lopezia coronata</i> ,	Coronet-flowered Lopezia, ..	3	220
<i>Lophospermum erubescens</i> ,..	Blushing Lophospermum, ..	5	401
<i>Lotus Jacobæus</i> ,	St. James's Island Trefoil, ..	4	326
<i>Lubinia atropurpurea</i> ,.....	Dark purple Lubinia,.....	5	461
<i>Lupinus Mexicanus</i> ,	Mexican Lupine,.....	4	366
<i>Lupinus mutabilis</i> ,	Changeable Lupine,	4	338
<i>Lupinus ornatus</i> ,	Adorned Lupine,.....	5	438
<i>Lupinus perennis</i> ,	Perennial Lupine,	4	310
<i>Lupinus polyphyllus</i> ,	Many-leaved Lupine,	4	291
<i>Lychnis Chalcedonica</i> ,	Double Scarlet Lychnis,....	1	87
<i>Lychnis coronata</i> ,	Chinese Lychnis,	2	148
<i>Lychnis flos Jovis</i> ,	Umbellate Rose-Campion, ..	5	397
<i>Lysimachia angustifolia</i> ,	Narrow-leaved Loose-strife, ..	2	104
<i>Lysimachia dubia</i> ,	Doubtful Loose-strife,.....	6	564
<i>Lythrum diffusum</i> ,	Diffuse Lythrum,.....	5	470
<i>Lythrum virgatum</i> ,	Fine-branched Lythrum,	2	134
<i>Macleaya cordata</i> ,	Heart-leaved Macleaya,.....	4	354
<i>Madia elegans</i> ,	Elegant Madia,	5	429
<i>Magnolia conspicua</i> ,	Yulan Magnolia,.....	3	236
<i>Malcomia maritima</i> ,	Mediterranean Stock,.....	3	196
<i>Malope trifida</i> ,.....	Trifid Malope,	3	281
<i>Malva angustifolia</i> ,.....	Narrow-leaved Mallow,	6	517
<i>Malva Mauritanica</i> ,	Mauritanian Mallow,	4	347
<i>Malva Munroana</i> ,	Munro's Mallow,	5	473
<i>Mathiola annua</i> ,	Ten-week Stock,	3	260
<i>Maurandia Barclayiana</i> ,	Barclay's Maurandia,.....	4	337
<i>Meconopsis Cambrica</i> ,.....	Welch Meconopsis,	5	479
<i>Melissa grandiflora</i> ,	Great-flowered Balm,.....	2	146
<i>Melittis melissophyllum</i> ,	Large-flowered Bastard Balm	4	372

<i>Systematic Name.</i>	<i>English Name.</i>	<i>Vol.</i>	<i>No.</i>
Melittis melissophyllum,....	Melissa-leav. Bastard-Balm,	4	356
Menziesia globularis,	Globe-flowered Menziesia,..	1	89
Menziesia polifolia,	Hoary-leaved Menziesia, ..	5	478
Mimulus luteus,	Yel.-flow. Monkey-flower,	2	116
Mimulus luteus, rivularis, ..	Spotted Yel. Monkey-flower,	4	308
Mimulus moschatus,	Musk Monkey-flower,	2	282
Mirabilis Jalapa,	Common Marvel of Peru, ..	1	16
Momordica elaterium,.....	Spirting Cucumber,	6	527
Monarda didyma,	Oswego Tea,	2	140
Monarda fistulosa,	Hollow-stalked Monarda, ..	6	571
Muscari comosum,	Tasselled Grape Hyacinth ..	3	250
Muscari comosum monstrosum,	Feathered Grape Hyacinth,	3	222
Narcissus angustifolius,	Narrow-leaved Narcissus, ..	5	407
Narcissus bicolor,	Two-coloured Narcissus,....	4	368
Narcissus bulbocodium,	Hoop Petticoat,	6	548
Narcissus incomparabilis, ..	Peerless Narcissus,	6	514
Narcissus interjectus,	Great curled-cup Narcissus,	5	412
Narcissus jonquilla,.....	Jonquil,	3	225
Narcissus tazetta,	Polyanthus Narcissus,	3	244
Neja gracilis,	Slender Neja,	5	393
Nicotiana tabacum,	Virginian Tobacco,.....	2	160
Nierembergia intermedia, ..	Nar-leav. pur. Nierembergia,	5	477
Nierembergia phœnicia,	Purple Nierembergia,	5	416
Nolana paradoxa,	Violet-flowered Nolana,....	3	207
Nuttallia papaver,	Poppy-flowered Nuttallia, ..	6	489
Œnothera densiflora,	Close-flowered Œnothera, ..	6	540
Œnothera Fraseri,	Fraser's Evening Primrose,	2	163
Œnothera Lindleyana,	Lindley's Tree Primrose, ..	3	201
Œnothera macrocarpa,	Large-fruited Œnothera, ..	1	41
Œnothera pumila,	Dwarf Œnothera,	1	35
Œnothera purpurea,	Purple-flowered Œnothera,	1	79
Œnothera roseo-alba,.....	Red and White Œnothera,..	4	365
Œnothera serrulata,	Saw-leav. Evening Primrose,	4	330
Œnothera speciosa,.....	Showy Œnothera,	4	339
Œnothera viminea,.....	Twiggy Œnothera,	4	362
Ononis hircina,	Strong-scented Rest-harrow,	2	174
Ononis rotundifolia,	Round-leaved Rest-harrow,	3	269
Onosma Taurica,.....	Golden-flowered Onosma, ..	3	234
Ophiopogon spicatus,	Spiked Snake's-Beard,	6	491
Opuntia vulgaris,	Common Opuntia,	6	552
Orobus hirsutus,	Hairy Bitter Vetch,	6	487
Orobus niger,	Black Bitter Vetch,	3	284
Orobus vernus,	Spring Bitter Vetch,	1	28
Oxalis crenata,	Notch-petalled Wood Sorrel,	5	433
Pæonia edulis,.....	Rose-scented Eatable Pæony,	5	451
Pæonia edulis, Whitleji,....	Eatable Pæony,	4	311
Pæonia moutan,	Poppy-flowered Tree Pæony,	3	241
Pancratium Illyricum,	Illyrian Pancratium,	2	159

<i>Systematic Name.</i>	<i>English Name.</i>	<i>Vol.</i>	<i>No.</i>
Pardanthus Chinensis,	Chinese Pardanthus,	6	481
Parnassia Caroliniana,	Carolina Parnassia,	6	550
Parnassia palustris,	Grass of Parnassus,	3	195
Passiflora cærulea,	Common Passion Flower,	1	4
Pentstemon atropurpureum ..	Dark Purple Pentstemon, ..	3	275
Pentstemon digitalis,	Foxglove-flow. Pentstemon, ..	3	258
Pentstemon ovatum,	Oval-leaved Pentstemon,	4	331
Pentstemon pubescens,	Hairy Pentstemon,	1	42
Pentstemon pulchellum,	Pretty Pentstemon,	4	335
Pentstemon Richardsonii,	Richardson's Pentstemon, ..	5	430
Pentstemon roseum,	Rose-coloured Pentstemon, ..	4	316
Pentstemon speciosum,	Showy Pentstemon,	4	373
Periploca Græca,	Grecian Periploca,	5	410
Petunia nyctaginiflora,	Large-flowered Petunia,	3	208
Phalangium liliago,	Grass-leaved Phalangium, ..	4	290
Phlomis herba-venti,	Wind Herb,	5	464
Phlomis tuberosa,	Tuberous Phlomis,	4	360
Phlox cæthea,	Fine-leaved Phlox,	1	68
Phlox crassifolia,	Thick-leaved Phlox,	4	319
Phlox divaricata,	Early blue-flowered Phlox, ..	2	182
Phlox nivalis,	Snow-white Phlox,	2	190
Phlox ovata,	Oval-leaved Phlox,	1	59
Phlox paniculata,	White-panicked Phlox,	4	350
Phlox pilosa,	Hairy-leaved Phlox,	4	309
Phlox reflexa,	Reflexed-leaved Phlox,	3	214
Phlox subulata,	Awl-leaved Phlox,	3	248
Phlox triflora,	Three-flowered Phlox,	1	6
Phlox Wheeleriana,	Wheeler's Phlox,	5	395
Phyteuma campanuloides,	Campanula-flow. Rampion, ..	2	158
Phyteuma orbiculare,	Round-headed Rampion,	2	154
Pinguicula grandiflora,	Large-Flowered Pinguicula, 6	560	
Piptanthus Nepalensis,	Nepal Piptanthus,	5	418
Polygala Chamæbuxus,	Box-leaved Milkwort,	1	24
Polygonatum multiflorum, ..	Many-flow. Solomon's Seal, ..	3	255
Polygonum divaricatum,	Divaricated Polygonum,	2	106
Polygonum viviparum,	Viviparous Polygonum,	3	282
Pontederia cærulea,	Blue Pontederia,	6	551
Potentilla atrosangui-pedata, ..	Hybrid Potentilla,	5	385
Potentilla formosa,	Handsome Nepaul Cinquefoil, 2	157	
Potentilla molissima,	Soft-leaved Cinquefoil,	6	542
Potentilla pedata,	Foot-leaved Cinquefoil,	4	379
Potentilla rupestris,	Rock Cinquefoil,	5	474
Potentilla Russelliana,	Russell's Hybrid Cinquefoil, ..	4	304
Potentilla splendens,	Splendid Cinquefoil,	3	221
Primula acaulis,	Double Primrose,	1	60
Primula auricula,	Auricula,	3	203
Primula auricula,	Auricula,	4	345
Primula cortusoides,	Cortusa-leaved Primrose,	2	156

<i>Systematic Name.</i>	<i>English Name.</i>	<i>Vol.</i>	<i>No.</i>
<i>Primula farinosa</i> ,	Bird's-eye Primrose,	1	96
<i>Primula minima</i> ,	Least Primrose.....	6	495
<i>Primula Sinensis</i> ,	Chinese Primrose,	1	1
<i>Primula villosa</i> , flore albo, ..	White Villous Auricula,	3	231
<i>Prunella grandiflora</i> ,	Great flowered Self-heal,....	2	102
<i>Prunella Sibirica</i> ,	Siberian Self-heal,	5	475
<i>Pulmonaria paniculata</i> ,	Panicled Longwort,.....	2	131
<i>Pulmonaria Virginica</i> ,	Virginian Longwort,	2	133
<i>Pyrethrum inodorum</i> ,	Scentless Feverfew,	4	359
<i>Pyrethrum roseum</i> ,	Rosy Feverfew,.....	4	380
<i>Pyrethrum uliginosum</i> ,.....	Marsh Feverfew,	5	439
<i>Pyrola minor</i> ,	Smaller Winter-green,	6	575
<i>Pyrus Japonica</i> ,	Japan Apple-tree,	1	49
<i>Ramonda Pyrenaica</i> ,	Borage-leaved Ramonda, ..	1	83
<i>Ranunculus amplexicaulis</i> , ..	Stem-clasping Ranunculus,..	4	321
<i>Ranunculus Asiaticus</i> ,	Garden Ranunculus,	3	271
<i>Ranunculus millefoliatus</i> ,....	Thousand-leaved Crowfoot,..	5	476
<i>Ranunculus platanifolius</i> ,....	Plane-tree-leaved Crowfoot, 4		295
<i>Rhexia Virginica</i> ,.....	Virginian Rhexia,	2	129
<i>Rhodanthe Manglesii</i> ,	Capt. Mangles's Rhodanthe, 6		537
<i>Rhodochiton volubile</i> ,	Twining Rodochiton,	6	525
<i>Rhododendron Catawbiense</i> ,..	Catawba Rhododendron,	5	408
<i>Rhododendron ferrugineum</i> ,..	Rusty-leaved Rhododendron, 4		325
<i>Rhododendron hirsutum</i> ,	Hairy-leaved Rhododendron, 1		63
<i>Rhododendron maximum</i> ,....	Large Rhododendron,	3	276
<i>Ribes aureum</i> ,	Golden-flowered Currant, ..	2	189
<i>Ribes sanguineum</i> ,	Crimson-flowered Currant, ..	4	377
<i>Rosa alba</i> ,.....	Celestial Rose,.....	6	561
<i>Rosa Banksiæ</i> , lutea,	Lady Banks's Yellow Rose,..	3	197
<i>Rosa centifolia</i> ,.....	Variegated Provins Rose,....	5	465
<i>Rosa centifolia</i> , var. Dianth.	Oeillet, or Pink-flow. Rose,..	3	249
<i>Rosa Indica</i> ,.....	China Rose,	2	185
<i>Rosa Indica</i> ,.....	Rose Clare,	5	445
<i>Rosa rapa</i> ,	Double Burnet-leaved Rose, 5		406
<i>Rosa semperflorens</i> ,	Crimson Chinese Rose,	2	177
<i>Rubus arcticus</i> ,.....	Dwarf Bramble,	2	152
<i>Rubus spectabilis</i> ,.....	Showy Bramble,	6	568
<i>Rudbeckia columnaris</i> ,.....	Columnar Rudbeckia,	4	336
<i>Rudbeckia fulgida</i> ,	Fulgid Rudbeckia,	2	101
<i>Rudbeckia hirta</i> ,	Hairy Rudbeckia,.....	4	314
<i>Rudbeckia pinnata</i> ,	Pinnate Rudbeckia,.....	5	457
<i>Sagittaria latifolia</i> ,	Broad-leaved Arrowhead, ..	6	546
<i>Salpiglossis straminea</i> ,.....	Straw-coloured Salpiglossis, 4		333
<i>Salvia grandiflora</i> ,	Great-flowered Sage,	5	420
<i>Sanguinaria Canadensis</i> ,	Bloodwort,	3	242
<i>Sanguisorba Canadensis</i> ,	Canadian Great Burnet,	5	442
<i>Saponaria Calabrica</i> ,	Calabrian Soapwort,.....	6	510
<i>Saponaria ocymoides</i> ,	Basil-leaved Soapwort,	2	651

<i>Systematic Name.</i>	<i>English Name.</i>	<i>Vol.</i>	<i>No.</i>
<i>Saxifraga aizoon</i> ,	Marginated Saxifrage,	4	419
<i>Saxifraga ceratophylla</i> ,	Shining-calyxed Saxifrage, 3	268	
<i>Saxifraga granulata</i> , fl. pleno, Dbl.	grain-root Saxifrage, ..	3	264
<i>Saxifraga oppositifolia</i> ,	Opposite-leaved Saxifrage, ..	2	127
<i>Saxifraga rotundifolia</i> ,	Round-leaved Saxifrage, ..	3	270
<i>Scabiosa atropurpurea</i> ,	Sweet Scabious,	2	119
<i>Schizanthus porrigens</i> ,	Spreading Schizanthus	2	126
<i>Schizanthus retusus</i> ,	Retuse-petalled Schizanthus, 6	521	
<i>Scilla bifolia</i> ,	Two-leaved Squill,	2	125
<i>Scilla bifolia alba</i> ,	White two-leaved Squill, ..	2	176
<i>Scilla Sibirica</i> ,	Siberian Squill,	2	192
<i>Scutellaria alpina</i> ,	Alpine Scullcap,	5	398
<i>Scutellaria columnæ</i> ,	Heart-leaved Scullcap,	2	141
<i>Scutellaria lupulina</i> ,	Two-coloured wolf Scullcap 6	555	
<i>Scutellaria peregrina</i> ,	Foreign Scullcap,	6	486
<i>Sedum aizoon</i> ,	Yellow Stonecrop,	4	348
<i>Sedum Ewersii</i> ,	Ewer's Stonecrop,	6	513
<i>Sedum oppositifolium</i> ,	Opposite-leaved Stonecrop, 5	460	
<i>Senecio elegans</i> ,	Purple Grounsel,	1	40
<i>Sida malvæflora</i> ,	Mallow-flowered Sida,	3	237
<i>Silene acaulis</i> ,	Stemless Catchfly,	3	232
<i>Silene fimbriata</i> ,	Fringe-flowered Catchfly, ..	2	128
<i>Silene laciniata</i> ,	Jagged-petalled Catchfly, ..	6	499
<i>Silene maritima</i> ,	Sea Catchfly,	4	324
<i>Sisyrinchium Bermudianum</i> ,	Bermuda Sisyrinchium,	4	381
<i>Sisyrinchium Bermudianum</i> ,	Large Bermuda Sisyrinchium, 6	497	
<i>Sisyrinchium grandiflorum</i> , ...	Large-flowered Sisyrinchium, 6	573	
<i>Sisyrinchium striatum</i> ,	Streaked Sisyrinchium,	1	66
<i>Solanum dulcamara</i> ,	Woody Nightshade,	2	109
<i>Solanum runcinatum</i> ,	Runcinate-leav. Nightshade, 6	520	
<i>Soldanella alpina</i> ,	Alpine Soldanella,	4	317
<i>Sollya heterophylla</i> ,	Various-leaved Sollya,	6	530
<i>Spigelia Marilandica</i> ,	Indian Pink,	1	93
<i>Spiræa bella</i> ,	Pretty Spiræa,	5	409
<i>Spiræa lævigata</i> ,	Smooth Spiræa,	1	32
<i>Spiræa trifoliata</i> ,	Three-leaved Spiræa,	2	153
<i>Statice oleæfolia</i> ,	Olive-leaved Sea Lavender, 1	86	
<i>Statice sinuata</i> ,	Sinuuous Sea Lavender,	6	500
<i>Stenactis speciosa</i> ,	Showy Stenactis,	6	547
<i>Stevia purpurea</i> ,	Purple Stevia,	2	180
<i>Symphoria, racemosa</i> ,	Snowberry,	2	117
<i>Symphytum Bohemicum</i> ,	Red-flowered Comfrey,	1	71
<i>Tagetes lucida</i> ,	Lucid Tagetes,	3	215
<i>Tagetes patula</i> ,	French Marigold,	1	56
<i>Tetragonolobus purpureus</i> , ..	Deep-red Tetragonolobus, ..	4	375
<i>Tetragonolobus siliquosus</i> , ..	Square-podded Winged Pea, 5	414	
<i>Teucrium fruticans</i> ,	Shrubby Germander,	6	554
<i>Teucrium Pyrenaicum</i> ,	Pyrenean Teucrium,	1	80

<i>Systematic Name.</i>	<i>English Name.</i>	<i>Vol.</i>	<i>No.</i>
<i>Thermopsis fabacea</i> ,.....	Fabia-like Thermopsis,	5	469
<i>Thymus lanuginosus</i> ,	Woolly Thyme,	2	184
<i>Tiarella cordifolia</i> ,	Heart-leaved Tiarella,.....	3	216
<i>Tigridia Pavonia</i> ,.....	Tiger Flower,	1	5
<i>Tolpis barbata</i> ,.....	Yellow Hawkweed,.....	3	259
<i>Tournefortia, heliotropioides</i> ,	Heliotrope-like Tournefortia,	6	483
<i>Trachymene cærulea</i> ,	Blue-flowered Trachymene,	5	450
<i>Tradescantia congesta</i> ,.....	Close-flowered Spider-wort,	4	363
<i>Trientalis Europæa</i> ,.....	European Winter-green,.....	6	544
<i>Trillium grandiflorum</i> ,.....	Large-flowered Trillium,	1	26
<i>Tritoma media</i> ,.....	Lesser Tritoma,	3	204
<i>Trollius Europæus</i> ,	European Globe-flower,	3	209
<i>Tropæolum majus</i> ,	Greater Nasturtium,	5	427
<i>Tulipa Gesneriana</i> ,	Common Tulip,	3	245
<i>Tulipa suaveolens</i> ,	Sweet-scented Tulip,	2	175
<i>Tulipa sylvestris</i> ,	Wild Tulip,	6	488
<i>Tussilago fragrans</i> ,	Sweet-scented Coltsfoot,	1	19
<i>Uvularia grandiflora</i> ,	Large-flowered Uvularia,	2	187
<i>Uvularia puberula</i> ,	Downy Uvularia,.....	6	512
<i>Vaccinium amœnum</i> ,	Broad-leaved Wortleberry,	3	211
<i>Valeriana montana</i> ,	Mountain Valerian,.....	1	36
<i>Verbascum phœniceum</i> ,	Purple-flowered Mullein,	1	36
<i>Verbena alata</i> ,.....	Winged-stemmed Vervain,	5	390
<i>Verbena Aubletia</i> ,	Rose-flowered Vervain,	3	205
<i>Verbena chamædrifolia</i> ,	Germander-leaved Vervain,	4	293
<i>Verbena pulchella</i> ,	Pretty Verbena,	3	277
<i>Verbena radicans</i> ,	Rooting Vervain,	5	422
<i>Verbena urticifolia</i> ,	Nettle-leaved Vervain,	5	488
<i>Veronica exaltata</i> ,	Lofty Speedwell,.....	6	543
<i>Veronica pinnata</i> ,.....	Wing-leaved Speedwell,.....	4	294
<i>Veronica urticæfolia</i> ,	Nettle-leaved Speedwell,	2	167
<i>Vesicaria utriculata</i> ,	Smooth Vesicaria,	1	84
<i>Vicia pseudocracca</i> ,.....	False Cracca Vetch,	6	507
<i>Vinca herbacea</i> ,	Herbaceous Periwinkle,	3	233
<i>Viola tricolor</i> ,	Heartsease,	5	421
<i>Viscaria neglecta</i> ,.....	Neglected Rock Lychnis,	6	523
<i>Wedelia aurea</i> ,.....	Golden-flowered Wedelia,	4	364
<i>Wistaria frutescens</i> ,	Frutescent Wistaria,	5	386
<i>Xeranthemum annuum</i> ,	Annual Xeranthemum,	2	114
<i>Xeranthemum lucidum</i> ,	Shining Xeranthemum,	2	135
<i>Yucca gloriosa</i> ,	Glorious Adam's-Needle,	3	286
<i>Zephyranthes Atamasco</i> ,	Atamasco Lily,	3	285
<i>Zexmenia tagetiflora</i> ,	Tagetes-flowered Zexmenia,	6	501
<i>Zinnia Elegans</i> ,.....	Elegant Zinnia,	6	549
<i>Zinnia elegans</i> ,.....	Purple-flowered Zinnia,	2	170
<i>Zinnia multiflora</i> ,	Red-flowered Zinnia,	2	97
<i>Zinnia tenuiflora</i> ,	Narrow-petalled Zinnia,	5	394
<i>Zizyphus paliurus</i> ,	Christ's Thorn,	2	118

PLATE I





Primula Sinensis

2/5



Amaryllis lutea

3/5



Iris versicolor

4/5



Passiflora carulea

1/5

PRIMULA SINENSIS.

CHINESE PRIMROSE.

Class.
PENTANDRIA.

Order.
MONOGYNIA.

Natural Order.
PRIMULACEÆ.

Native of China.	Height. 6 inches.	Flowers in Mar. July.	Duration. Perennial.	Introduced in 1822.
---------------------	----------------------	--------------------------	-------------------------	------------------------

No. 1.

Primula is derived from the Latin primus, first, from its early flowering; hence its English name also, prime-rose, now contracted to primrose. Sinensis, from Sinæ, the name of an ancient people, who are supposed to have inhabited that part of the Chinese empire now called Cochin China.

This plant is the Primula prænitens of the Botanical Register; but as the term Sinensis has not been established for another species, upon authority worth naming, we give it the preference, on account of its prior adoption.

The attention of the Horticultural Society was first drawn to this beautiful plant in the year 1819, when a drawing of it was received from John Reeves, Esq., a corresponding member, residing at Canton. A plant, and seeds also, were subsequently sent off by him to the Society; the former perished during its passage, and the latter did not vegetate. Since that period it has been introduced by Capt. Rawes, and as it possesses beauties so completely distinct from every other primula with which we are acquainted, it is likely to become a distinguished favourite in our gardens.

Its mode of inflorescence is particularly beautiful; for out of a simple umbel or head of flowers, rises a distinct scape or stalk, supporting a second umbel, and from this is produced a third, and sometimes a fourth; by which peculiarity, and its free increase of flowering side shoots, it remains in bloom during the greater part of the year. Our drawing was taken at an early stage of the blossom, and represents the first umbel only.

Having been introduced so lately, its habits are but imperfectly known; it may, however, be considered hardy, as plants of it were exposed in the open air during the winter of 1823, in different parts of the kingdom, without sustaining injury.

It may be propagated very readily by offsets, and flourishes exceedingly in a pot of compost, made with equal parts of peat, rich loam, and sand; or it may be planted in a warm, dry border, of light soil, and have the protection of a hand-glass during severe frost.

Seeds are produced very freely by the *Primula Sinensis*, and from them young plants may be propagated in abundance, and with little trouble. They should be sown as early as March, in pots of light rich earth, placed in a hotbed; and the young plants, when large enough, should be potted singly, and be gradually inured to the open air; but they will require occasional shade in the summer. The protection of the cold frame is necessary for them during the first winter of their growth, and in April, part may be removed to dry parts of the borders, for flowering; and part may be retained in pots, as portable summer or winter ornaments.

Lindley's Col. Bot.

AMARYLLIS LUTEA.

YELLOW AMARYLLIS.

Class.
HEXANDRIA.

Order.
MONOGYNIA.

Natural Order.
AMARYLLIDÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
Spain.	4 inches.	Sept. Oct.	Perennial.	in 1596.

No. 2.

Amaryllis, the name of a shepherdess in Theocritus and Virgil. Lutea, yellow, its colour. It is sometimes called Autumnal Narcissus.

Known as this hardy, fast increasing, plant has been for two or three hundred years, it is remarkable that it should not be nearly as common as our yellow crocus, to which, at first sight, it appears so closely allied. But it is not the harbinger of spring : it does not excite the delightful sensations which every daisy, every buttercup of that joyous season, is calculated to arouse.

“What lovely prospects wait each wakening hour,
When each new day some novelty displays;
How sweet the sunbeam melts the crocus flower,
Whose borrow'd pride shines dizen'd in his rays.” CLARE.

It grows well in almost any soil or situation, except under the dripping of trees ; for as its bulbs are reproduced but slowly, during the severity of winter, the leaves continue to increase till spring ; at which time the bulbs will become fully matured, and the leaves will die. Transplanting may then be performed with propriety, till vegetation is reassumed in July.

Aiton's Hort. Kew. ed. 2, vol. 2. p. 223.

AMARYLLIS LUTEA

YELLOW AMARYLLIS

Class. *Liliaceae*
Order. *Liliales*
Suborder. *Liliales*
Family. *Liliaceae*
Genus. *Amaryllis*
Species. *A. lutea*

Native of	Spain
Height	4 inches
Flowers in	April, Oct.
Duration	Perennial
Published	in 1808

Amaryllis, the name of a subgenus in *Lilium*, is given to *Lilium* and *Veril*. *Lilium*, yellow, the color. It is sometimes called *Amaryllis*. Known as this name, but is common, it is common for two or three hundred years. It is common that it should not be easily as common as our yellow *Lilium*, to which, at first sight, it appears as closely allied. But it is not the harbinger of spring. It does not excite the delightful sensations which every day, every hour of that joyful season, is calculated to arouse.

It grows well in almost any soil or situation, except under the shelter of trees, where its bulbs are exposed, but always, during the season of winter, the leaves continue to increase till spring, at which time the bulbs will be unusually enlarged and the leaves will die. Transplanting may then be performed with impunity, till vegetation is resumed in July. *Amaryllis lutea*, L. p. 222.

IRIS VERSICO'LOR.

CHANGEABLE IRIS.

Class.
TRIANDRIA.

Order.
MONOGYNIA.

Natural Order.
IRIDEÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
N. America	15 inches.	June.	Perennial.	in 1732.

No. 3.

This plant has received its name from the Latin term Iris, which signifies a rainbow; and the colours of some of the species render it very appropriate. It has been termed the various-coloured, or parti-coloured; changeable-coloured appears a more correct translation, and is equally characteristic of the flower, for it may be observed daily to assume a different hue.

An eastern, or even northern, border is suitable to this plant. It flourishes in any light garden soil, and the roots may be divided in autumn. It may also be raised from seeds, which should be sown in September, and the plants will come up in the following spring; but if the seeds are sown in the spring, they will lie a year in the ground before they vegetate.

That correct observer of nature, Bradley, speaking of one of the bulbous Irises, says, the finest varieties that he ever saw, were raised from seed; 'I would,' he further observes, 'advise every one to raise seedlings.'

Orris root is the tuber of the Florentine Iris, which will hereafter be noticed.

Hort. Kew. 2, v. 1, 116.

THIS VERSICOLOR.

CHANGING IRIS.

Color
Horizontal

Color
Horizontal

Vertical Color
Horizontal

Plant in N. America in 1850	Height in inches	Flowers in June	Flowers in August	Cultivated in 1731
--------------------------------	---------------------	--------------------	----------------------	-----------------------

No. 3

This plant has received its name from the Latin term *iris*, which signifies a rainbow; and the colors of some of the species render it very appropriate. It has been termed the various-colored, or parti-colored; changeable-colored; appears a more correct translation, and is equally characteristic of the flower, for it may be observed daily to assume a different hue.

An eastern, or even northern, border is suitable to this plant. It thrives in any light garden soil, and the seeds may be divided in autumn. It may also be raised from seeds, which should be sown in September, and the plants will come up in the following spring; but if the seeds are sown in the spring, they will bear fruit in the ground before they vegetate.

That correct observer of nature, Baskley, speaks of one of the Indian tribes, says, the first variety that he ever saw, were raised from seed; 'I would,' he further observes, 'advise every one to raise seedlings.'

Our root is the tuber of the *Peruvian Iris*, which will better be named.

From the *Bot. & L. Lib.*

PASSIFLORA CÆRULEA.

COMMON PASSION FLOWER.

Class.

PENTANDRIA.*

Order.

TRIGYNIA.

Natural Order.

PASSIFLOREÆ.

Native of Brasil.	Height. 30 feet.	Flowers in Aug. Oct.	Duration. Perennial.	Cultivated in 1699.
----------------------	---------------------	-------------------------	-------------------------	------------------------

No. 4.

Passiflora is derived from the Latin *patior*, to suffer; and *flos*, a flower; from the fancied resemblance of the different parts of the flower and plant to the instruments of Christ's suffering. The five stamens were compared to his five wounds; the three styles, more aptly, to the nails by which he was fixed to the cross; the column which elevates the germen, to the cross itself; and the rays of the nectary, to his crown of thorns; the petals to the ten apostles, Judas and Peter being rejected; the tendrils to a cord, the leaf to a hand, &c. &c. *Cærulea*, from the Latin, blue.

Parkinson, in his *Paradisus Terrestris*, gives what he calls 'The Iesuite's Figure of the Marococ,' which is a representation of the flower, composed of the very instruments of torture themselves; but in noticing these fancies, he is very angry at the superstition that suggested them; observing that it is 'All as true as the sea burnes.'

It may be propagated from seeds, cuttings, or

* Sir J. E. Smith, in his excellent Introduction to Physiological Botany, coincides with Schreber and Thunberg, in placing this genus in the class Pentandria.

layers. Cuttings may be taken early in the spring, of the preceding year's growth ; or in June, of the young shoots, and struck under a hand-glass. It sometimes ripens its seed in the open air in England ; and these may be sown in pots, placed in a hotbed, in March ; and the plants gradually exposed to the open air after Midsummer.


Miller says, 'I have found the plants which have been propagated two or three times, either by layers or cuttings, seldom produce fruit ; which is common to many other plants.' This observation of Miller's should not be lost sight of. It may, in many instances of fruit culture, prove very important, as the mode of propagation has undoubted influence on the habits of many vegetables.

Parkinson, in his notice of the maracoc, which was a species of the passion flower, cultivated when he wrote in 1629, says, that it showed a remarkable particularity in rising from the ground a month sooner, if a seedling plant, than if it grew from roots brought from Virginia.

It appears highly desirable to propagate from seeds, which has been done by several eminent botanists, and beautiful new varieties have been produced. Some of these are minutely described in the Transactions of the Horticultural Society.

The *Passiflora cærulea* requires to be trained against a wall, with a southern aspect ; and in winter the roots should be covered with straw, as this will protect them from injury, even if the head of the plant be destroyed. In March the shoots may be very much shortened, which will encourage a vigorous growth.

Hort. Kew. 2, v. 4, 154.





Tigridia Pavonia

$\frac{1}{3}$



Phlox triflora

$\frac{1}{2}$



Helleborus niger

$\frac{1}{2}$



Fragaria Indica

$\frac{1}{2}$

TIGRIDIA PAVONIA.

TIGER FLOWER.

Class.

MONANDRIA.

Order.

TRIANDRIA.

Natural Order.

IRIDEE.

Native of Mexico.	Height. 1 foot.	Flowers in July, Sep.	Duration. Perennial.	Cultivated in 1796.
----------------------	--------------------	--------------------------	-------------------------	------------------------

No. 5.

Ferraria Pavonia, Linneus's name of the present plant, was derived from John Baptista Ferrarius, who first figured and described the *Ferraria undulata*. Linneus made choice of his specific name, *Pavonia*, in consequence of Mutis having sent him a drawing of the flower under this appellation, which he had given it from Pavon, the name of a favourite pupil; and not, as is generally supposed, from the Latin pavo, a peacock, on account of the beauty of its colours.

This species is now separated from *Ferraria*; and its present generic name, *Tigridia*, is deduced from tigris, a tiger, by reason of its spots. *Leopardia* would, perhaps, on such account, have been a more appropriate appellation.

The exquisite union of colours, and conformation of parts, in this beautiful production, abash every attempt of the pencil; description or portraiture can but do it discredit. We regret its visit being so transient, opening in the morning and finally closing in the afternoon; and yet it is certain that much of our pleasure depends on such circumstances. Sturm justly observes, "If flowers retained their beauty

throughout the year, they would not impart to us the delight they now do: their absence makes us long for their return. The constant variation and succession of all terrestrial objects, constitute one of the chief sources of our happiness."

If the bulbs be planted in the borders, or on separate beds, about the middle of April, the spring frosts will have ceased before the leaves appear through the soil, and no protection be required to be given them. To produce earlier flowers, they may be put into pots, in a common hotbed, in the beginning of March, and watered sparingly till the leaf appears. If a sufficiency of air be allowed them, they will bear exposure early in May; and then should be turned into the borders for flowering, taking care to retain the balls of earth quite perfect about the roots, which will be much assisted by a copious watering supplied a few hours before their removal. They neither increase as fast, nor flower quite as freely, if left in the pots. The roots should be taken up before frosts commence, and those bulbs which are attached together should remain so; then be gradually dried, and afterwards secured in paper bags, and kept in a cool dry place till spring.

The *Tigridia* increases its bulbs rapidly, and also produces seeds, from which it may be raised by sowing them in pots, in a hotbed, in the spring. The seedling bulbs will, of course, require taking up as before directed; and they will flower, some in the second, and the remainder in the third year.

The root is an ovate bulb, which is eatable when roasted, tasting like a chestnut.

PHLOX TRIFLORA.

THREE-FLOWERED PHLOX.

Class.
PENTANDRIA.

Order.
MONOGYNIA.

Natural Order.
POLEMONIACEÆ.

Native of N. America.	Height. 2 feet.	Flowers in August.	Duration. Perennial.	Introduced in 1815.
--------------------------	--------------------	-----------------------	-------------------------	------------------------

No. 6.

Linneus did not, in this genus of plants, retain the old name *Lychnidea*, in consequence of its near approach to *Lychnis*, the name of another genus; but chose the present name *Phlox*, perhaps, only by reason of its analogy to the former, both being derived from Greek words, signifying flame or lightning. *Triflora*, from the branches of the corymbus being mostly three-flowered.

Every individual of this family inherits some desirable quality; the greater part of them are extremely showy; several grow higher than the present plant; and a few clothe the border with a close foliage during the whole year. England now possesses about thirty species of *Phlox*, brought principally from North America, within the last eighty or ninety years.

It will grow in any common soil, but best in a mixture of peat and loam. It may be easily increased by cuttings, taken early in the summer, or by dividing the roots, in autumn; but this latter practice should not be resorted to oftener than once in two or three years, or the roots will be much weakened.

Sweet's Fl. Gar. p. 29.

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

THEORY OF THE

FRAGA'RIA IN'DICA.

YELLOW-FLOWERED STRAWBERRY.

Class.
ICOSANDRIA.

Order.
POLYGYNIA.

Natural Order.
ROSACEÆ.

Native of E. Indies.	Height. 4 inches.	Flowers in May, July.	Duration. Perennial.	Introduced in 1804.
-------------------------	----------------------	--------------------------	-------------------------	------------------------

No. 7.

Fragaria, so named from the fragrancy of its fruit. Indica, from its native country.

The English name, Strawberry, is, without doubt, derived from the practice of laying straw about the roots of such plants, to preserve the fruit from the soil. Straw, slates, &c. used in this way yield two advantages;—at the same time as the fruit is kept clean, the soil is preserved in a moist and cool state.

It is principally remarkable for its union of the cinquefoil blossom with the fruit of the strawberry, and certainly forms a pretty variety amongst the closer sorts of rock plants.

The *Fragaria Indica* is frequently treated as a greenhouse plant, but is now found to bear our winters without injury. Its fruit is of no further value than for its ornamental appearance, not possessing the prominent characteristics of its tribe, fragrance and flavour.

It propagates itself readily by its emission of roots from the joints, as the common varieties usually do; and flourishes in a sandy soil, where the situation is warm and sheltered.

Hort. Kew. 2, v. 3, 273.

FRAGARIA INDICA

YELLOW-FLOWERED STRAWBERRY.

Order.
FRUITFUL.

Class.
FRUITFUL.

Native of the
North.

Native of the North.	Native of the North.	Native of the North.	Native of the North.	Native of the North.
Native of the North.	Native of the North.	Native of the North.	Native of the North.	Native of the North.

No. 7.

Fraser, as named from the frequency of its fruit, is a native country.

The English name, Strawberry, is, without doubt, derived from the practice of laying straw about the roots of each plant, to preserve the fruit from the soil. Straw, straw, &c. used in this way yield two meanings;—at the same time as the fruit is kept warm, the soil is preserved in a moist and cool state.

It is principally remarkable for its union of the compound character with the fruit of the strawberry, and certainly forms a variety among the other sorts of fruit plants.

The Fraser's Indica is frequently treated as a greenhouse plant, but is now found to bear our winter without injury. The fruit is of no further value than for its ornamental appearance, not possessing the prominent characteristics of its tribe, fragrance and flavor.

It propagates itself freely by its emission of roots from the joints, as the common varieties usually do; and thrives in a sandy soil, where the situation is warm and sheltered.

Herb. Kew. t. 2. 573.

HELLEBORUS NIGER.

CHRISTMAS ROSE.

Class.
POLYANDRIA.

Order.
POLYGYNIA.

Natural Order.
RANUNCULACEÆ.

Native of Austria.	Height. 9 inches.	Flowers in January.	Duration. Perennial.	Cultivated in 1596.
-----------------------	----------------------	------------------------	-------------------------	------------------------

No. 8.

Helleborus, from the Greek, expressive of its poisonous qualities; or, according to Bergeret, from the river Eleborus. Niger, from the Latin, black, the external colour of its root.

The flower, at its first opening, is white, afterwards rather pink, and finally becomes green. The tubular nectaries ranged round the germen, merit the attention of the physiologist.


A tincture of the roots of black hellebore, is employed in medicine; but as its effects are somewhat uncertain and dangerous, we forbear giving its mode of preparation. It may not be amiss, however, for the information of those who use the roots, to observe that others, and sometimes still more dangerous ones, are substituted for them. The following description of the genuine is from the Edinburgh Dispensatory.

“The roots consist of a black furrowed roundish head, about the size of a nutmeg, from which short articulated branches arise, sending out numerous corrugated fibres, about the thickness of a straw, from a span to a foot in length, deep brown on the outside, white or yellowish white within, and of an

acid, nauseous, and bitterish taste, exciting a sense of heat and numbness in the tongue, and of a nauseous acid smell. These fibres only are used in medicine, and the head and decayed parts are rejected. For the roots of the real black hellebore, the roots of the *Adonis vernalis*, *Trollius Europæus*, *Actæa spicata*, *Astrantia major*, *Helleborus viridis foetidus*, *Veratrum album*, and *Aconitum neomontanum* are often substituted. The last is a most virulent poison, and may be distinguished by its roots being fusiform, or nearly globular, sending out numerous very brittle fibres, of a greyish black or brown colour, as thick as a man's finger, and repeatedly divided."

If the virtues of this plant, like those of many others, were formerly too much extolled, they are, probably, now undeservedly neglected; it is indeed to be regretted that the study of medical botany has, of late years, made so little progress; or, it may rather be said, that in the last age it should have so far declined. Modern chemists are, however, showing the potent effects of condensed vegetable properties, which promises fair to constitute a new era in medical knowledge.

In a moist situation, where a little peat has been mixed with the soil, this plant flowers abundantly; and the flower stems will grow higher, and the blossoms be altogether improved, by the assistance of a hand glass, which may be placed over them on four small pots, so as to admit a free current of air underneath. It is readily increased by dividing the roots, which is best effected in the early part of autumn.





Globularia vulgaris.

$\frac{1}{2}$



Cuscuta verrucosa.

$\frac{1}{2}$



Erinus alpinus.

$\frac{1}{2}$



Jasminum revolutum.

$\frac{1}{2}$

GLOBULARIA VULGARIS.

COMMON GLOBULARIA.

Class.
TETRANDRIA.

Order.
MONOGYNIA.

Natural Order.
GLOBULARIÆ.

Native of Europe.	Height. 6 inches.	Flowers in May, June.	Duration. Perennial.	Cultivated in 1629.
----------------------	----------------------	--------------------------	-------------------------	------------------------

No. 9.

This plant was named by Tournefort, from the flowers growing many together, in the form of a little globe or ball. Parkinson says "The Italians call it Botanaria, because the heads are found like buttons."

The *Globularia vulgaris* is a pretty close-growing plant, which was classed by several of our old botanists with the garden daisy; and called *Bellis cærulea*, or blue daisy. In some situations it seldom blossoms so freely as may be wished; but this defect will generally be found to arise from its situation being too dry and warm: hence it is well adapted to a northern border, or the cool side of artificial rock wo.

It may be propagated from seeds, or by parting the roots, which is best effected in September, when the plants will have an opportunity of making new roots before frosts commence. It flourishes in a shady situation, in a light rich soil or in sandy peat; and to encourage a good bloom, should not be too frequently transplanted. It will succeed very well if kept in a pot, and placed, during winter, in a cold frame, with the alpine plants.

CUSCUTA VERRUCOSA.

WART-CALYXED DODDER.

Class.
PENTANDRIA.

Order.
DIGYNIA.

Natural Order.
CONVOLVULACEÆ.

Native of	Height.	Flowers in	Duration.	Introduced
Nepaul.	6 feet.	September.	Perennial.	in 1822.

No. 10.

Some authors have deduced the word *Cuscuta*, from the modern Greek, others from a similar Asiatic word. *Verrucosa*, from the Latin *verruca*, a wart. The English appellation, *Dodder*, seems to have been derived from the German word *dotter*, or Dutch *tauteren*, signifying to shoot up.

This is a plant which twines round ivy or shrubs of any description that it comes in contact with; and though raised from seeds in the soil, as are most other vegetables, yet no sooner does it meet with support from a neighbouring branch, than, like many individuals of a superior order of creation, it quits its original friend and supporter, and clings to a new acquaintance. It twines in a direction contrary to the apparent course of the sun, and throwing out little vesicles, which attach themselves to the plant that supports it, thereby draws its necessary nutriment, and dies off at the root, becoming completely parasitical.

It flourishes most on soft succulent shrubs; and should severe frosts destroy it, young plants may be raised from seed in the spring; and they will produce their fragrant little flowers in autumn.

Sweet's Fl. Gar. 6.

ERSCUTA VERMICOLA

WART-CAYEN WORM

Order
DIOPTERA

Family
TENTHEDINIDAE

General Description
of the Larva

Length of Larva	Length of Larva	Length of Larva
1.5 mm.	1.5 mm.	1.5 mm.

No. 10

Some authors have chosen the word *Coccus* from the modern Greek, others from a similar Latin word, *Vermis*, from the Latin *vermis*, a worm. The English appellations, *Worm*, *Worm*, *Worm*, have been derived from the German word *Worm*, or Dutch *worm*, signifying a worm.

This is a plant which grows round the roots of shrubs of any description that it comes in contact with; and though it is found on the roots, as we have seen, it does not grow there, but it feeds on the roots of other vegetables, and on roots that it meets with support from the surrounding earth, like many individuals of a species of *Veronica*, it feeds on the roots of shrubs and vegetables, and also on a new acquisition. It is very common in the garden, and the apparent cause of the loss, and throwing out little roots, which often themselves to the plant that supports it, thereby shows its necessary nature, and this is at the root, becoming completely parasitic.

It furnishes most on soft succulent shrubs; and should never be destroyed, if young plants are to be raised from seed in the spring; and they will give their first growth like flowers in autumn.

Worm's Life.

ERINUS ALPINUS.

ALPINE ERINUS.

Class.
DIDYNAMIA.

Order.
ANGIOSPERMIA.

Natural Order.
SCROPHULARINEÆ.

Native of Pyrenees.	Height. 3 inches.	Flowers in April, May.	Duration. Perennial.	Cultivated in 1759.
------------------------	----------------------	---------------------------	-------------------------	------------------------

No. 11.

The Greek term ERINOS, whence comes our Erinus, was applied by the ancients to a vegetable, very different from any contained in the present genus; it was the Wild Fig of the Greeks, and its name was, probably, derived from a verb, implying to exert or strive, because the Greek plant endeavoured to erect itself by means of walls or stones. Alpinus, from the Latin, belonging to the Alps.

This interesting little subject presents us with its pretty flowers at that season, when all animated nature seems most capable of such enjoyment.

“ Propitious spring comes forth in bright array,
With Venus, goddess of the vernal day;
Her mild precursor, Zephyr, wafts the breeze,
With balmy wings, o’er all the budding trees:
Maternal Flora, with benignant hand.
Her flowers profusely scatters o’er the land.
These deck the vallies with unnumber’d hues,
And far around their fragrant sweets diffuse.”

The Erinus alpinus requires a dry shady situation. On building rubbish, or in the open joints of an old wall, it succeeds admirably. It may be propagated by seeds, or by division.

Hort. Kew. 2, v. 4, 49.

ERINUS ALPINE.

ALPINE ERINUS.

Order.
Labiata.

Class.
Dicotyled.

General Order.
Natural Order.

Height of the plant.	Height of the flower.	Height of the fruit.	Height of the seed.
-------------------------	--------------------------	-------------------------	------------------------

Pl. 11.

The Greek name *Erinus*, whence comes our *Erinus*, was applied by the ancients to a vegetable, very different from any contained in the present genus; it was the *Urtica* of the Greeks, and its name was, probably, derived from a verb, signifying to exert or strive, because the Greek plant endeavored to erect itself by means of walls or stems, *Alpinus*, from the Latin, belonging to the Alps.

This interesting little subject presents us with its pretty flowers at that season, when all animated nature seems most capable of such enjoyment.

"Erinus, which comes forth in bright array,
With green, green leaves of the velvet dye;
Her mild perfume, *Euphry*, wafts the breeze,
While beauty weeps on all the bounding trees;
Her flowers, I love, with fragrant hand,
Her flowers, I love, with fragrant hand,
Then deck the walls with numerous'd flowers,
And far around their fragrant sweets diffuse."

The *Erinus alpinus* requires a dry shady situation. On building rubbish, or in the open joints of an old wall, it succeeds admirably. It may be propagated by seeds, or by division.

Herb. Kew. v. 4. 40.

JASMINUM REVOLUTUM.

CURLED-FLOWER JASMINE.

Class.
DIANDRIA.

Order.
MONOGYNIA.

Natural Order.

JASMINÆ.

Native of	Height.	Flowers in	Duration.	Introduced
China.	16 feet.	May, Aug.	Perennial.	in 1812.

No. 12.

Jasminum is derived from two Greek words, signifying a violet, and odor, on account of the fine scent which its flowers possess. Revolutum, in allusion to the curling of its petals.

We are told by Capt. Hardwicke, in the Asiatic Researches, that a species of yellow Jasmine, which is supposed to be the present one, was observed by him, on the frontiers of Hindostan, and which formed a large bush. We have never seen the Jasminum revolutum treated as a standard shrub, though it appears well calculated for that purpose, if sufficiently hardy. Trained against a southern wall, it puts forth vigorous shoots, and produces abundance of flowers of the richest fragrance, and most brilliant golden hue.

It may be propagated readily by cuttings, taken in the spring, and appears to grow very well in any common soil. A little straw or matting should be laid over the roots to protect them from frosts of long continuance; and in very inclement seasons, a slight covering over the branches, also will prove beneficial, by preserving the young and succulent shoots, which the luxuriant habit of the plant will leave rather too


tender to withstand the severity of our winters, though the older branches remain uninjured.

The flowers of this, as well as the common Jasmine, are admirably adapted to the use of the toilette ; and to some of our fair readers the following method of extracting perfumes, from the produce of their gardens, may not be wholly unacceptable. We copy it from the Family Receipt Book.

‘ Procure a quantity of the petals of any flowers which have an agreeable fragrance ; card thin layers of cotton, which dip into the finest Florence or Lucca oil ; sprinkle a small quantity of fine salt on the flowers, and lay them, a layer of cotton and a layer of flowers, until an earthen vessel or a wide-mouthed glass bottle is full. Tie the top close with a bladder, then lay the vessel in a south aspect to the heat of the sun, and in fifteen days, when uncovered, a fragrant oil may be squeezed away from the whole mass, little inferior, if that flower is made use of, to the dear and highly valued Otto or Odor of Roses.’

When the aroma or odor is united with a rectified spirit, it is usually termed an essence ; and this may readily be obtained by mixing a portion of the oil, prepared as above directed, with an equal quantity of alcohol ; shake them together in a phial, and the spirit will become impregnated with the perfume of the oil. They may afterwards be poured from each other, and the essence preserved for use.

From the *Jasminum officinale* the Italians obtain an essential oil, by distillation, which is held in high estimation amongst them as a remedy for rheumatic pains, and for application to paralytic limbs.





Coreopsis tinctoria

$\frac{7}{8}$



Linum alpinum

$\frac{3}{2}$



Catananche cœrulea

$\frac{7}{8}$



Mirabilis jalapa

$\frac{3}{2}$

COREOPSIS TINCTORIA.

ARKANSA COREOPSIS.

Class.
SYNGENESIA.

Order.
POLYGAMIA FRUSTRANEA.

Natural Order.
CORYMBIFERÆ.

Native of N. America.	Height. 3 feet.	Flowers in Aug. Oct.	Duration. Annual.	Introduced in 1823.
--------------------------	--------------------	-------------------------	----------------------	------------------------

No. 13.

Coreopsis, from two Greek words, signifying the appearance of a bug or tick, in allusion to the fancied resemblance of its seed to such insects. Tinctoria, from the Latin tinctura, a colour or dye, expressive of the colouring properties of its petals.

The appellation Arkansa, indicates the situation, in North America, where it was discovered by Professor Nuttall, during his travels through that country. Nearly all the individuals belonging to the genus Coreopsis, are interesting hardy subjects, chiefly natives of the same inexhaustible and comparatively unexplored, source of the wonders of nature.

The Coreopsis tinctoria is a remarkably pretty slender-growing annual, and having been lately introduced amongst us, is by no means common. It will flourish in any rich soil, and may be treated as are the usual annuals of our gardens, by being sown at the end of March. If early flowers are desired, the seed may be sown a fortnight sooner, in a hotbed, and the plants removed into the borders in April. In dry situations autumn-sown plants will frequently endure the winter.

LINUM ALPINUM.

ALPINE FLAX.

Class.

PENTANDRIA.

Order.

PENTAGYNIA.

Natural Order.

CARYOPHILLÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
Austria.	8 inches.	June, Aug.	Perennial.	in 1739.

No. 14.

Linum comes from the Greek word LINON, signifying flax or cotton; or cloth manufactured from either of them. Alpinum, from the Latin alpinus, belonging to the Alps. Our English word flax, is of Saxon origin, from flex.

This little plant resembles the Linum usitatissimum, or common flax, so well known; but is shorter in the stem, and the flowers are much larger.

It may be propagated by cuttings, which strike readily under a hand-glass; or the seeds may be sown as soon as they are ripe. It is a pretty plant for rock work or the fronts of borders, and prefers a dry sandy soil.

Flax appears to have been cultivated by the Egyptians, upwards of three thousand years ago, for we read of it in the book of Exodus, chap. 9, ver. 31; though it does not appear at what precise period it first became employed for the purposes to which we now convert it.

Notwithstanding the mention of linen is found in the works of the most ancient historians, it is not quite certain that it was, in their time, manufactured either of flax or hemp.

The Greeks are said, at a very early period, to have been supplied with their fine linen from Egypt; but their term LINON applies indiscriminately to cloth of cotton or flax; and it is probable, that the former was used many ages before the latter.

Herodotus, the Greek historian, who lived about two thousand years ago, and who travelled in Egypt, frequently notices the use of linen, but we neither find it determined by the historian himself, nor any of his commentators, that flax was, at that period, manufactured into cloth. It may have been cultivated for its seed only.

Linseed oil, which is expressed from the seed of the common flax, is highly valuable to the arts, particularly as a component part of paint.

An infusion of flax seed has been found to be extremely useful as a pectoral drink, in coughs, colds, and affections of the lungs. It is thus prepared:—Take an ounce of bruised flax seed, half an ounce of sliced liquorice root, and pour upon them a quart of boiling water; let it macerate an hour, near the fire, and then strain it off for use. A teacup-full may be taken five or six times in the course of twenty-four hours; when a fresh supply should be made.

Equal parts of linseed oil and lime water, form a liniment that no family should be without. It is singularly useful in the cure of burns and scalds; and if timely applied, prevents the inflammation attendant on these cases.

The flax stem when macerated in streams or ponds, is said to communicate a poisonous quality to the water; and an act of Henry VIII. prohibited the practice, under pain of twenty shillings.

Hort. Kew. 2, v. 2, 187.

CATANANCHE CÆRULEA.

BLUE-FLOWERED CATANANCHE.

Class.
SYNGENESIA.

Order.
POLYGAMIA ÆQUALIS.

Natural Order.
CICHORACEÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
S. Europe.	2 feet.	July, Sep.	Perennial.	in 1596.

No. 15.

Catananche is derived from two Greek words, signifying compulsion; from its supposed power of causing an irresistible impulse to love; a quality which, formerly, it was ridiculously imagined to possess. Cærulea, from the Latin, blue.

The blue-flowered Catananche appears to have been an old inhabitant of our gardens, and it is said to be synonymous with the *sesamoides parvum* of Gerarde; if so, the figure given of it by that author, is certainly inferior to most of those with which he has supplied us.

Miller observes that it may be propagated by heads taken from the mother plant. This we have never tried, finding it easily raised from seeds; and we conceive, with more success than by dividing.

Sow seeds of the Catananche cærulea in March or April, in rich light soil, and when the young plants come up, they should be thinned, if required, and kept clear from weeds till autumn. The seedlings may then be planted out, one in a place, in dry situations, where they are intended to remain, and an abundance of flowers will be produced in the following summer.

Hort. Kew. 2, v. 4, 469.

CATAPACHIA GARRINER

1875, BENTLEY AND CATAPACHIA

On the 1st of May, 1875, the following was received from the Hon. the Secretary of the Navy, Washington, D.C.:

Specimens of Catapachia	Specimens of Catapachia	Specimens of Catapachia	Specimens of Catapachia
1875, May 1st	1875, May 1st	1875, May 1st	1875, May 1st

No. 12

Catapachia is derived from two Greek words, signifying compulsion; from its supposed power of causing an irresistible impulse to love; a quality which, however, it was erroneously imagined to possess. Catapachia, from the Latin, plus.

The blue-flowered Catapachia appears to have been an old inhabitant of our gardens, and it is said to be synonymous with the semiothisis pattern of the garden; it is, the figure given of it by that author, is certainly inferior to most of those which he has supplied us.

Miller observes that it may be propagated by seeds taken from the mother plant. This we have never tried, finding it easily raised from seeds; and we conceive, with more success than by dividing.

Some seeds of the Catapachia were in March set apart in rich light soil, and when the young plants came up, they should be thinned, if required, and kept clear from weeds till autumn. The seedlings may then be planted out, one in a place, in dry situations, where they are intended to remain, and an abundance of flowers will be produced in the following summer.

That No. 12, 1875.

MIRA'BILIS JALA'PA.

COMMON MARVEL OF PERU.

Class.

PENTANDRIA.

Order.

MONOGYNIA.

Natural Order.

NYCTAGINEÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
India.	3 feet.	August.	Perennial.	in 1596.

No. 16.

This plant is said to have received its name, *Mirabilis*, from the 'wonderful diversity of colours in the flowers.' We may venture an opinion that it arose from the diverse combination of them; for the colours in a single flower are by no means numerous. One plant will produce many flowers entirely red, some of a clear yellow, and others variegated in different degrees, with red, yellow, and occasionally with cream colour. There is a purple and white variety also, which possesses the same changeful propensity in the disposition of its two colours, but we are not aware of these colours ever being mingled with the red and yellow; and seedling plants most frequently produce plain flowers only.

The specific name, *Jalapa*, was adopted on the supposition that the officinal Jalap was produced by the *Mirabilis*. This error has been corrected by Dr. Houston, who discovered, in the Spanish West Indies, that the plant from which the Jalap of the shops is prepared, is a species of *convolvulus*, and is now known as the *Convolvulus Jalapa*.

Bullock, when travelling in Mexico, in 1823, observed that Jalap was chiefly produced in the


neighbourhood of Xalapa: hence its appellation; the J and the X, in the Spanish language, having the same sound, and being interchangeable.

Our old herbalists, two centuries ago, were well acquainted with this plant, and 'dwell with great marvel thereon.' Parkinson calls it the 'Meruaile of the World;' and mentions, amongst others, one that he possessed with blossoms of a pale purple or peach colour.

It has obtained the appellation of the four-o'clock plant, from the flowers usually opening about that time in the afternoon. In cool or gloomy weather, however, they continue expanded during the next day; otherwise the warmth of the sun, early in the morning, closes them to open no more: when evening again arrives, we find our plant with fresh embellishments, as gay as before.

The Marvel of Peru is usually cultivated as a half-hardy annual; but it is far better to take up the roots on the first approach of frost, and lay them in dry sand, in a cellar, till the last week in March; when they may be replanted in the borders for flowering. As the roots, when thus put out into the borders, will sometimes remain in the ground a considerable time before they vegetate, it is preferable to plant them in deep pots of soil, and place them in a hotbed till they have grown an inch or two: they may then be turned out, with the balls of soil unbroken, into the situations in which they are intended to flower. If seedlings are desired, raise them in a hotbed, as early as is convenient, that they may blow before the sharp autumnal nights prevent it.

Hort. Kew. 2, v. 1, 382.





Gaultheria procumbens.

$\frac{3}{2}$



Amsonia latifolia.

$\frac{3}{2}$



Tussilago fragrans

$\frac{3}{2}$



Andromeda calyculata latifolia.

$\frac{3}{2}$

GAULTHERIA PROCUMBENS.

TRAILING GAULTHERIA.

Class.
DECANDRIA.

Order.
MONOGYNIA.

Natural Order.
ERICÆ.

Native of N. America.	Height. 5 inches.	Flowers in July, Aug.	Duration. Perennial.	Cultivated in 1768.
--------------------------	----------------------	--------------------------	-------------------------	------------------------

No. 17.

The present genus has been named after Dr. Gaulthier, or Gautier, a physician and botanist of Canada; whence the *Gaultheria procumbens* was first introduced. *Procumbens*, from the Latin, implying a bending downwards.

Our present is an interesting little subject, and is, in appearance, somewhat like the *vaccinium*, or bilberry tribe. Its pendent crimson berries sometimes remain on the plant and accompany the flowers of the following year. They form a pretty contrast with each other; and with its motley evergreen foliage, constitute a small, though brilliant, ornament in a shady part of the garden.

The inhabitants of some parts of North America are said to use the leaves as we do that incomparable produce of the 'Celestial Empire'—Tea.

An infusion of them certainly possesses considerable fragrance; and were we not prejudiced in favour of our accustomed beverage, might be thought equally pleasant; and would, probably, be found equally wholesome.

It should be planted in peat, in a cool situation; where it will slowly increase by its creeping roots.

AMSONIA LATIFOLIA.

BROAD-LEAVED AMSONIA.

Class.
PENTANDRIA.

Order.
MONOGYNIA.

Natural Order.
APOCYNÆ.

Native of N. America.	Height. 15 inches.	Flowers in June, July.	Duration. Perennial.	Cultivated in 1759.
--------------------------	-----------------------	---------------------------	-------------------------	------------------------

No. 18.

The appellation, Amsonia, appears to have been first given to this plant by the author of the Flora Virginica, Mr. Clayton, who discovered it in Virginia; but we are not told in what its name originated; a supposition may be ventured that he made choice of it from the name of some person or place. Latifolia, from the Latin latus, broad, and folium, a leaf.

This plant has long been an inhabitant of the English parterre, and is generally thought to have been cultivated by Miller. This, however, is not quite certain; for, in his dictionary, he speaks of the Tabernæmontana Amsonia, which is considered synonymous with the present Amsonia latifolia, as having white flowers; ours it will be seen are light blue.

It does not increase rapidly, though it may occasionally be divided at the root. It succeeds best in a warm situation where the soil is continually kept in a moderate degree of moisture. It may be planted in a compost of peat and loam, with the addition of a small quantity of sand, and should not be frequently removed.

Hort. Kew. 2, v. 2, 72.

AMORPHIA LATIFOLIA

BROAD-LEAVED AMORPH

White
Missouri

Common
Virginia

Western Canada
Atlantic

Height	15 to 20 feet
Flowers	White, July to August
W. America	in 1858

20 16

The supposition, *Amorpha*, appears to have been first given to this plant by the author of the *Flora Virginica*. Mr. Clayton, who discovered it in Virginia, but we are not told in what its name might be; a supposition may be ventured that he made it from the name of some person or place. Lathrop, from the Latin *latus*, broad, and *folium*, a leaf.

This plant has long been an inhabitant of the English prairie, and is generally thought to have been introduced by Miller. This, however, is not quite correct; for, in his description, he speaks of the *Tephrosia virginiana*, which is considered synonymous with the present *Amorpha latifolia*, as having white flowers; ours it will be seen are light blue.

It does not increase rapidly, though it may occasionally be divided at the root. It requires but a warm situation where the soil is occasionally kept in a moderate degree of moisture. It may be planted in a row of four and four, with the addition of a small quantity of sand, and should not be frequently removed.

Bar. Rec. 2, 73

TUSSILA'GO FRA'GRANS.

SWEET-SCENTED COLTSFOOT.

Class.

SYNGENESIA.

Order.

POLYGAMIA SUPERFLUA.

Natural Order.

CORYMBIFEREÆ.

Native of Italy.	Height. 8 inches.	Flowers in January.	Duration. Perennial.	Introduced in 1806.
---------------------	----------------------	------------------------	-------------------------	------------------------

No. 19.

The term, Tussilago, has been derived from the Latin tussis, a cough; and ago, to drive away; on account of the efficacy of one of its species in affections of the lungs. Fragrans, from the Latin, fragrant. The English name Coltsfoot has arisen from the shape of its leaves.

The bloom is odoriferous in a high degree, very similar to the Heliotropium Peruvianum; and if the plant be made an inhabitant of the greenhouse, or sitting room, when in flower, very few that we are acquainted with, afford so rich a perfume; though the peculiar almond-like scent may not be equally agreeable to all.

The foliage of our present subject, and its habits also, bear a strong resemblance to the Tussilago farfara, a native of our own fields; and with which many a worthy farmer regrets his too intimate acquaintance. It is, indeed, a rank weed, betraying a sterile soil.

Notwithstanding the Tussilago fragrans is highly valued for the odour of its blossoms, produced at a dreary period of the year; yet we warn the unwary florist against introducing it into his borders,

without first inclosing its roots within the narrow precincts of an earthen flower pot. If this be neglected his fragrant friend may, probably, by its creeping, or rather running, roots, make a rapid tour over the principal part of his garden.

Having noticed the *Tussilago farfara*, or common Coltsfoot, we may, perhaps, be allowed the present opportunity of making further mention of it.

Curtis, in his *Flora Londinensis* mentions a practice prevalent amongst the Tartars of carrying burning touchwood, which was probably made of the roots of *Tussilago*, the smoke being intended to protect them from the annoyance of gnats. Thus the invention of a pipe may have arisen in the necessity of sometimes employing the breath to quicken the fire.

He further observes, 'The custom of smoking this plant, which still prevails, is of ancient date. Pliny directs the dried leaves and root of Coltsfoot to be burnt, and the smoke drawn into the mouth through a reed and swallowed, as a remedy for an obstinate cough.'

Coughs have frequently been much relieved by persons smoking the dried leaves; and they are used as the principal ingredient in the manufacture of British Herb Tobacco. An infusion of the dried flowers, sweetened with honey, has been found particularly efficacious in complaints of this nature.

The downy substance which is found growing on the under surface of the leaves of the common coltsfoot, when dipped in a strong solution of saltpetre, and gradually dried, is said to form an admirable tinder.

Hort. Kew. 2, v. 5, 35.

ANDROMEDA CALYCVLATA. *latifolia.*

BROAD-LEAVED CALYCVLED ANDROMEDA.

Class.

DECANDRIA.

Order.

MONOGYNIA.

Natural Order.

ERICÆ.

Native of	Height.	Flowers in	Duration.	Introduced
Newfoundl ^d	2 feet.	Mar. Apr.	Perennial.	in 1748.

No. 20.

Andromeda, a celebrated classic beauty of antiquity, the daughter of Cepheus and Cassiope. She was bound by the nymphs to a rock, to be devoured by a sea monster, because her mother proudly preferred her beauty to theirs; but Perseus, rescued and married her. Calyculata, double calyxed, from the presence of two minute ovate leaflets, which grow on the base of the calyx.

This pretty evergreen shrub, with some trifling variations in its growth, is found native nearly all round the northern parts of the globe; as in Siberia, Sweden, and North America. Being highly astringent to the taste, its medicinal qualities are probably of a tonic nature. The combined circumstances of situation and properties in the Andromeda calyculata are such as M. Pastié of the Royal Academy of Sciences of Paris may advance in support of his new hypothesis. He thinks it of the highest importance, in making ourselves acquainted with the medicinal properties of plants, that we attend to their native situations; and even conceives this of greater consequence than knowing to what genus they belong, or the chemical principles which

enter into their composition. He observes, that all plants which grow on high cold grounds have a tonic and stimulant power; whilst those are found of contrary qualities which are natives of opposite situations.

“Nor ev’ry plant on ev’ry soil will grow:

The sallow loves the wat’ry ground, and low;

The marshes, alders; Nature seems t’ordain

The rocky cliff for the wild ash’s reign:

The baleful yew to northern blasts assigns,

To shores the myrtles, and to mounts the vines.”

DRYDEN’S VIRGIL.

Santolius Victorinus, a French writer, seems to have had the above lines of Virgil in mind when he wrote the following; which occur in Evlyn’s translation of M. Quintinye’s French Gardener, a work showing much experience, combined with considerable ingenuity and talent.

“All soils affect not every sort of stock,

The apple chooseth earth, the pear the rock;

The peach flies marshes, some delight to share

The hottest sun, and choose an open air.


Some love the shade, here trees and shrubs will spread,

There flowers from seed adorn a noble bed.”

The severest cold will not affect the *Andromeda calyculata*, but it is desirable with this, as with most other of the northern plants, that they be kept somewhat shaded during the continuance of our hot months.

It should be planted in a mixture of peat and fresh loam; and may be propagated by layers, which will not be sufficiently rooted for separation in less time than two years.

Hort. Kew. 2, v. 3, 55.





Lachenalia tricolor

7/8



Erica herbacea



Lonicera Tartarica

7/2



Polygala Chamæbuxus

LACHENALIA TRICOLOR.

THREE-COLOURED LACHENALIA.

Class.

HEXANDRIA.

Order.

MONOGYNIA.

Natural Order.

ASPHODELEÆ.

Native of	Height.	Flowers in	Duration.	Introduced
C. G. Hope.	8 inches.	April, May.	Perennial.	in 1774.

No. 21.

The term *Lachenalia* was derived from the name of Wernerius de la Chenal, an eminent botanist of Switzerland. *Tricolor*, three coloured, from the presence of three colours on the flowers, but these have very little permanency. They change considerably as the flowering advances.

Many bulbous plants possess a singular mode of reproduction, in the formation of little buds or bulbs on their flowering stems, which admit of separation; and, being properly treated, grow to perfection. The bulb-bearing lily, (*lilium bulbiferum*,) and tiger lily, (*lilium tigrinum*,) are familiar examples of this peculiarity.

A still more singular occurrence is stated by Sir J. E. Smith; who says that he has had scaly bulbs form even on the flower stalk of the *Lachenalia tricolor*, whilst lying for many weeks between papers to dry; which, on being put into the ground, became perfect plants.

The production of a viviparous progeny, under such circumstances, is well calculated to remind us that no means are too difficult to be employed by the Almighty for the preservation of an individual

even of the lowest order of creation. The farther we explore the operations of nature, the more shall we find to delight and surprise us.

Nothing can exceed the ingenuity of the various contrivances, if we may be allowed these expressions, that exists for the more certain propagation of many of our commonest plants. Seeds appear to be the legitimate source of reproduction in vegetables; but nature admits no bounds to her efforts. Where her usual purposes meet with opposition, she employs other means for the accomplishment of the same ends. Numerous plants produce seeds very abundantly, and at the same time a vigorous production takes place at the roots; whilst others increase very slowly by the latter means. We may, however, in general, observe that the vegetating power of herbaceous plants, if not required for the seasonable support of seeds, will be exerted in the increase of their species, either by the multiplication of tubers or bulbs, beneath or above the soil; by suckers, by runners, offsets, or by some other mode, agreeably to the nature of the plant. These observations cannot, of course, apply to exotics growing in a soil or climate uncongenial to their nature.

The *Lachenalia tricolor* will succeed in a warm border, with the protection of a hand glass in severe frosts. Or, which is frequently more convenient, it may be planted in a pot of light soil, composed of equal parts of fine sand, rotten leaves, and fresh loam, and be afforded the protection of a cold frame, or other sheltered situation during frost; and such treatment will insure success. It should be very sparingly watered during winter.

Hort. Kew. 2, v. 2, 288.

ERICA HERBA'CEA.

EARLY DWARF HEATH.

Class.
OCTANDRIA.

Order.
MONOGYNIA.

Natural Order.
ERICÆ.

Native of Austria.	Height. 6 inches.	Flowers in Jan. Mar.	Duration. Perennial.	Introduced in 1763.
-----------------------	----------------------	-------------------------	-------------------------	------------------------

No. 22.

From the Greek EREICO, to break, from its supposed quality of breaking, or rather dissolving, the stone in the bladder. Herbacea, from the Latin, herbaceous. It is sometimes called Erica carnea. Several other synonyms also have been applied to this species, but they are not in use.

Perhaps no tribe of plants yields more ample evidence of the rapid progress of floriculture in this country than the present.

Miller, not sixty years ago, described but five sorts of heath as known in England; but at the present period, there are upwards of three hundred distinct species. Out of this number, according to Dr. Withering's arrangement, five only are indigenous to Great Britain; and, unfortunately, very few of the exotics will bear the severity of our climate, having been chiefly introduced from the Cape of Good Hope.

This heath may be raised from cuttings, but as its procumbent branches increase freely, in any light soil, a convenient mode of propagation is offered by layers, which will root sufficiently to admit of separation in eighteen months.

Hort. Kew. 2, v. 2, 366.

LONICE'RA TARTA'RICA.

TARTARIAN HONEYSUCKLE.

Class.
PENTANDRIA.

Order.
MONOGYNIA.

Natural Order.
CAPRIFOLEÆ.

Native of Russia.	Height. 5 feet.	Flowers in Apr. May.	Duration. Perennial.	Cultivated in 1752.
----------------------	--------------------	-------------------------	-------------------------	------------------------

No. 23.

Adam Lonicer, in honour of whom this genus has been named, was a physician and botanist of Frankfurt, where he died in 1588. Tartarica, from Tartary, whence seeds were sent to Petersburg, and thence to England.

This species of *Lonicera* constitutes a pretty variety amongst low-growing shrubs. Its delicate spring foliage contrasts admirably with the deep green tints of Portugal laurel, *lauristinus*, and others of our favourite evergreens.

In its native country it is said to be frequented by the *Lytta vesicatoria*, or blistering fly. They are gathered from it by shaking the branches over a cloth, and after being killed by the fumes of vinegar, are dried in a stove, and preserved for use. These are sometimes mixed with another insect, the *melontha vitis*, which does not stimulate the skin. The latter is easily distinguished by its shape, which is nearly square, and should be rejected as useless for the purposes intended.

This *Lonicera* is of easy propagation, either by layers or cuttings, but does not generally succeed in the smoke of towns.

Hort. Kew. 2, v. 1, 379.

LONICERA TARTARICA TARTARIAN HONEY-SUCKLE

SYDNEY
UNIVERSITY

LIBRARY
HERBARIUM

RECEIVED
JAN 21 1884

Native of	Height	Flowers in	Duration	Collected
India	5 feet	Apr. May	Perennial	in 1882

No. 21

Adam Lonicer, in honour of whom this genus has been named, was a physician and botanist of Frankfort, where he died in 1588. Tartarian, from Tartary, whence seeds were sent to Peterburgh, and thence to England.

This species of Lonicera constitutes a pretty variety amongst low-growing shrubs. Its delicate spring foliage contrasts admirably with the deep green tints of Portugal laurel, laurustinus, and others of our favourite evergreens.

In its native country it is said to be propagated by the Lytta vesicatoria, or blistering fly. They are gathered from it by shaking the branches over a cloth, and after being killed by the fumes of vinegar, are dried in a stove, and preserved for use. These are sometimes mixed with another insect, the mallochia vicia, which does not stimulate the skin. The latter is easily distinguished by its shape, which is nearly square, and should be rejected as useless for the purposes intended.

This Lonicera is of easy propagation, either by cuttings or suckers, but does not generally succeed in the method of sowing.

Herb. Kew. E. 1, 377.
Bot. Beech. E. 1, 377.

POLY'GALA CHAMÆBUX'US.

BOX-LEAVED MILKWORT.

Class.
DIADELPHIA.

Order.
OCTANDRIA.

Natural Order.
PEDICULAREÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
Austria.	6 inches.	Apr. May.	Perennial.	in 1658.

No. 24.

This genus appears to have obtained its name from the supposition that some of its species increase the quantity of milk in cows that feed upon it. It is a compound of two Greek words, POLU, much, and GALA, milk. By some of our botanists of the seventeenth century, the present subject was called Chamæbuxus, from the Greek, signifying low-growing box; hence the term has been retained as a trivial name.

This is a most desirable low evergreen shrubby plant, producing a profusion of fragrant flowers, during the months of April and May; and partially through the summer.

Several varieties of the Polygala chamæbuxus are spoken of; some with red flowers, others with red and yellow; but we believe they are not at present known in England.

It grows extremely well if planted in peat, or peat and loam, and increases freely by its creeping roots. When a removal or division of the plant is required, it may be safely effected in April, or the beginning of September, care being taken to water it if requisite.

Hort. Kew. 2, v. 4, 245.

POLY-GALA CHAMBERLAIN

BOX-LEAVED HILLWORT.

Class.
Diptera

Order.
Hymenoptera

Suborder.
Hymenoptera

Number of Antennae	Length of Body	Length of Wing	Length of Foreleg	Length of Hindleg
10	1.5	1.5	1.5	1.5

No. 22


This genus appears to have obtained its name from the supposition that some of its species increase the quantity of milk in cows that feed upon it. It is a compound of two Greek words, *polu*, much, and *gala*, milk. By some of our botanists of the seventeenth century, the present subject was called *Chamberlain*, from the Greek, signifying low-grow-ing box; hence the term has been retained as a tri-
vial name.

This is a most desirable low evergreen shrubby plant, producing a profusion of fragrant flowers, during the months of April and May; and partially through the summer.

Several varieties of the *Polygala chamberlain* are spoken of; some with red flowers, others with red and yellow; but we believe they are not at present known in England.

It grows extremely well if planted in peat, or peat and loam, and increases freely by its creeping roots. When a removal or division of the plant is required, it may be easily effected in April, or the beginning of September, care being taken to water it if repotted.

Bot. Rev. 2, v. 4, 182.





Dodecatheon Meadia.

$\frac{1}{2}$



Trillium grandiflorum.

$\frac{1}{2}$



Adonis vernalis.

$\frac{2}{3}$



Orobus vernus.

$\frac{1}{2}$

DODECA'THEON MEAD'IA.

AMERICAN COWSLIP.

Class.

PENTANDRIA.

Order.

MONOGYNIA.

Natural Order.

PRIMULACEÆ.

Native of Virginia.	Height. 12 inches.	Flowers in May.	Duration. Perennial.	Introduced in 1744.
------------------------	-----------------------	--------------------	-------------------------	------------------------

No. 25.

Dodecatheon is formed of two Greek words, signifying twelve gods. Meadia was the only name this plant first received, and it was given to it by Catesby, in honour of Dr. Mead, an English physician of considerable eminence. It is not quite obvious why Linneus rejected Meadia, and adopted Dodecatheon, as a generic term; a name applied by Pliny, it is supposed, to our English cowslip. Meadia, as it stood named by Catesby, must be considered as reasonable an appellation, since Dr. Mead appears to have been a man fully worthy of the honour at first done him.

Very few plants excite more general interest than is produced by the Dodecatheon Meadia. It is one of those attractive flowers that will bear the most scrupulous examination, and still leave us the more in admiration of its beauties.

The grains of the farina or dust of this flower, when inspected with the assistance of a compound microscope, will be found to be peculiarly beautiful. They are distinctly organized minute pearls. So minute, that one square inch will contain of them upwards of three millions; and as squares cannot

be covered by circles, more than one fifth of the space will still be left unoccupied. Or, to be more particular in numbers, presuming that a square inch will contain three millions of circles, in direct rows each way, the area of each of such circles will be the 3,819,709th part of the area of an inch.

Mr. Phillips's simile, in his amusing work, the *Flora Historica*, is an apt one. He says the petals are reflexed, or turn back over the calyx, giving the appearance of an half-expanded parasol; a resemblance which is considerably heightened by the long tapering shape of the parts of fructification, and the golden colour of the anthers.

The flowers, during their expansion, assume an elegantly pendent position, as best adapted to their protection and fertilization; but when the farina has fallen, and the flower fades, the peduncles, with the seed vessels, become perfectly erect.

The chief care necessary in the cultivation of the *Dodecatheon*, is the planting it in a shady situation. It flourishes in any cool light soil, particularly in peat; and may be safely removed in the autumn, or early in the spring. It frequently ripens its seed, and from these may be readily propagated. If they are sown in pots in the autumn, they will vegetate in the spring, when great care should be taken that the young plants be not injured by the heat of the sun. Keep them in a shady situation, moderately moist, and clean from weeds. After their leaves are decayed, they may be removed, but this should not be deferred later than the middle of September; when they may be put into the borders to remain for flowering.

Hort. Kew. 2, v. 1, 311.

TRILLIUM GRANDIFLO'RUM.

LARGE-FLOWERED TRILLIUM.

Class.
HEXANDRIA.

Order.
TRIGYNIA.

Natural Order.
SMILACEÆ.

Native of N. America.	Height. 6 inches.	Flowers in April.	Duration. Perennial.	Cultivated in 1796.
--------------------------	----------------------	----------------------	-------------------------	------------------------

No. 26.

Trillium, from the triple formation of the several parts of the plant and flower. It is composed of three leaves, three stigmas, three petals, three calyx-leaves, and the berry is three-celled. Grandiflorum, from the Latin grandis, great; and flos, a flower. This plant has also been called Trillium erythrocarpum, red-fruited.

All the Trilliums, with which we are at present acquainted, are natives of America; growing there in woods and shady places. They may be raised from seeds, sown in September or October, which will come up early in the following spring. The young plants should be kept moist and cool during the summer, and in September, the roots may be transplanted into shady borders of light earth where they are intended to remain. In England, these plants increase very slowly at the root, and the original ones are frequently lost, even under the care of the most experienced cultivators; and if the seeds are sown in the spring, they remain twelve months dormant, which may lessen the chances of success. These combined circumstances tend to limit the whole of the Trilliums to the borders of the curious.

Par. Lond. t. 1.

and to TRIBULUM GRANDIFOLIUM.

grown on CLIFF-FLOWERS-VEGETATION.

It is a small plant, growing in the same places as the other species of the genus. It is a small plant, growing in the same places as the other species of the genus. It is a small plant, growing in the same places as the other species of the genus.

Height of plant	1 foot
Height of flower	1 foot
Height of fruit	1 foot
Height of seed	1 foot

The fruit is a small, round, black, and is very hard.

TRIBULUM, from the Latin tribulus, a thorn, or a prickly plant. It is a small plant, growing in the same places as the other species of the genus. It is a small plant, growing in the same places as the other species of the genus. It is a small plant, growing in the same places as the other species of the genus.

All the TRIBULUM, with which we are acquainted, are natives of America; growing there in woods and shady places. They may be raised from seeds, sown in September or October, which will come up early in the following spring. The young plants should be kept moist and cool during the summer, and in September, the roots may be transplanted into shady borders of light earth where they are intended to remain. In England, these plants increase very slowly at the root, and the original ones are frequently lost, even under the care of the most experienced cultivators; and if the seeds are sown in the spring, they remain twelve months dormant, which may lessen the chance of success. These combined circumstances tend to limit the whole of the TRIBULUM to the borders of the garden.

ADONIS VERNALIS.

SPRING ADONIS.

Class.
POLYANDRIA.

Order.
POLYGYNIA.

Natural Order.
RANUNCULACEÆ.

Native of Europe.	Height. 12 inches.	Flowers in Mar. April.	Duration. Perennial.	Cultivated in 1731.
----------------------	-----------------------	---------------------------	-------------------------	------------------------

No. 27.

Adonis, whom this flower has been named after, was the favourite of Venus. Vernalis, from the Latin, belonging to the spring.

Some of our fabulists have a pleasant conceit respecting these 'personages:' we relate it as being in some degree explicable. They say, that when Venus and Proserpina contended before Jupiter, which should have Adonis, Jupiter referred them to Calliope, whom he appointed to be judge of their quarrel. Calliope gave this sentence; that Adonis should live with Venus six months, and the remaining six he should reside with Proserpina. The fable is thus explained: Venus represents the earth, and Adonis the sun. During half the year his resplendent beams reign over all our varied plains, attired with beauteous flowers, and enriched with fruit and corn; the other half he seems to lose his influence, and goes as it were to rule in the darker regions with Proserpina.

To encourage a good blossom, the Adonis vernalis should not be frequently removed. If it be performed in the autumn, the flowering of the following spring may, possibly, receive but little injury.

Hort. Kew. 2, v. 3, 350.

ADONIS VERNALIS.

SPRING ADONIS.

Order.
POLYGYRIA.

Class.
POLYGYRIA.

Natural Order.
RANUNCULACEAE.

Native of Europe.	Height. 12 inches.	Flowers in Mar. April.	Domestic. Perennial.	Cultivated in 1731.
----------------------	-----------------------	---------------------------	-------------------------	------------------------

No. 27.

Adonis, whom this flower has been named after, was the favourite of Venus, Vernalis, from the Latin, belonging to the spring. Some of our fabulists have a pleasant conceit respecting these personages; we relate it as being in some degree applicable. They say, that when Venus and Proserpina contended before Jupiter, which should have Adonis, Jupiter referred them to Calliope, whom he appointed to be judge of their quarrel. Calliope gave this sentence; that Adonis should live with Venus six months, and the remaining six he should reside with Proserpina. The first is thus explained: Venus represents the earth, and Adonis the sun. During half the year his resplendent beams reign over all our varied plains, adorned with fragrant flowers, and enriched with fruit and corn; the other half he seems to lose his influence, and goes as it were to rule in the dark regions with Proserpina. To encourage a good husband, the Adonis vernalis should not be frequently removed. It is best sown in the autumn, the flowering of the following spring may possibly receive but little injury.

Hort. Kew. 2, t. 3, 350.

O'ROBUS VER'NUS.

SPRING BITTER VETCH.

Class.
DIADELPHIA.

Order.
DECANDRIA.

Natural Order.
LEGUMINOSÆ.

Native of Europe.	Height. 12 inches.	Flowers in April.	Duration. Perennial.	Cultivated in 1629.
----------------------	-----------------------	----------------------	-------------------------	------------------------

No. 28.

Orobus comes from the Greek oro, to excite; and BOUS, an ox; in allusion to the utility of some of the plants belonging to this genus in fattening cattle. Vernus, from the Latin ver, spring, its time of flowering.

This is the Orobus venetus of Gerarde and Parkinson, though the latter mentions it as double the height of our plant. He says, 'This pretty kinde of pease blossome beareth diuers slender, but upright, greene branches, somewhat cornered, two foote high, or thereabouts.'

It is, indeed, a very pretty kind of pease blossom, and so early a visiter cannot fail of being welcome in our borders. Its colour varies as the flowering advances; till, on fading, it becomes a light blue; and the gay diversity thus afforded, yields a pleasing effect.

We never have seen it produce any seeds. It may be divided for increase, in autumn, but its deeply penetrating strong roots should not be too frequently removed. A shady situation is usually recommended for the Orobus vernus; we find it, however, succeed in almost any soil or aspect.

Hort. Kew. 2, v. 4, 303.

O-ROBUS VERNUS.
SPRING BITTER VETCH.

Order.
DIPSACALES.

Class.
DICOTYLEDONES.

Natural Order.
LABIATAE.

Native of Europe.	Height. 12 inches.	Flowers in April.	Position of Pedicels.	Configuration of Leaves.
----------------------	-----------------------	----------------------	-----------------------------	--------------------------------

No. 32.

Orobos comes from the Greek *oros*, to excite; and *ros*, an ox; in allusion to the ability of some of the plants belonging to this genus in fattening cattle. *Vernus*, from the Latin *ver*, spring, its time of flowering.

This is the *Orobos vernus* of Gerard and Parkinson, though the latter mentions it as double the height of our plant. He says, 'This pretty kind of pease blossom bears a slender, but upright, green branch, somewhat curved, two foot high, or thereabouts.'

It is, indeed, a very pretty kind of pease blossom, and so early a visitor cannot fail of being welcome in our borders. Its colour varies as the flowering advances; till, on fading, it becomes a light blue; and the gay diversity thus afforded, yields a pleasing effect.

We never have seen it produce any seeds. It may be divided for pasture, in autumn, but its deeply penetrating strong roots should not be too frequently removed. A shady situation is usually recommended for the *Orobos vernus*; we find it, however, succeed in almost any soil or aspect.

Bot. Kew. 2. v. 4. 303.





Delphinium grandiflorum.

½



Iris Susiana.

¼



Lilium bulbiferum.

½



Spiraea laevigata.

DELPHINIUM GRANDIFLO' RUM.

LARGE-FLOWERED LARKSPUR.

Class.
POLYANDRIA.

Order.
TRIGYNIA.

Natural Order.
RANUNCULACEÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
Siberia.	2 feet.	June, Aug.	Perennial.	in 1758.

No. 29.

The appellation applied to this genus has proved less mutable, under the hand of science, than many others, though we are strongly inclined to believe that it has usurped the name of another tribe. The delphinium or Greek DELPHINION of Dioscorides, received its name from the fancied resemblance of its flower bud to a dolphin. Any one who will compare the unopened flowers of the several species of delphinium and aconitum, will readily observe, that the latter bear a much stronger similitude than the former, to that fish.

The beautiful double variety of Delphinium grandiflorum, so well known, and generally called the Siberian Larkspur, is an offspring of the plant before us; and is increased by division.

The present one may be increased by dividing the roots in the usual season; or by seeds, which are freely produced. They may be sown either in autumn or spring, but those sown at the former time will produce the strongest plants. These should be kept thin, and free from weeds, during the summer; and, not later than October, be transplanted into their proper situations for flowering.

DELPHINIUM GRANDIFLORUM

LARGE-FLOWERED LARKSPUR

Height	Flowers in	Flowers in	Flowers in
2 feet	June, Aug.	June, Aug.	June, Aug.
2 feet	June, Aug.	June, Aug.	June, Aug.

The appellation applied to this genus has proved less suitable, under the hand of science, than many others, though we are strongly inclined to believe that it has usurped the name of another genus. The delphinium or Greek variety of *Delphinium* is derived its name from the Greek resemblance of its flower bud to a dolphin. Any one who will examine the different flowers of the several species of *delphinium* and *acanthus*, will readily observe, that the latter bears a much stronger resemblance than the former, to that fish.

The beautiful double variety of *Delphinium* given to us, is well known, and generally called the *Delphinium*, is an offspring of the plant just mentioned, and is increased by division.

The question one may be increased by division, the roots in the usual season; or by seeds, which are very numerous. They may be sown either in autumn or spring, but those sown at the former time will produce the strongest plants. These should be kept thin, and free from weeds, during the summer; and not later than October, be transplanted into their proper situation for flowering.

Lib. R. N. 2, 360

Iris susiana L.

CHALCEDONIAN IRIS.

Class. TRIANDRIA. *Order.* MONOGYNIA.

Natural Order. IRIDACEÆ.

Native of Levant.	Height. 2 feet.	Flowers in May.	Duration. Perennial.	Cultivated in 1596.
----------------------	--------------------	--------------------	-------------------------	------------------------

No. 30.

Peculiar circumstances, or qualities, belonging to plants, which sometimes give rise to their generic names, will rarely be found equally applicable to all the individuals which must necessarily be included in the same genus. The Latin term *iris*, a rainbow, applies admirably to many of the plants bearing such name, but certainly, not to the present one, which has no such variety of colour.

Wonderful as were the exertions and penetration of the great Linneus, and eminently skilled as are many of his successors, still the efforts of science are inadequate, when applied to the developement of the laws of nature. Her laws are fixed, but so diversified, so complex, so utterly inexplicable to human understanding, that man, the boasted lord of the creation, must stand abashed by his ignorance, and science herself confess her defects.

Naturalists have zealously, and meritoriously, explored her mysteries, and endeavoured to assign to her, specific laws, whereby to circumscribe her economy; but still before the code could ever be completed, her numerous exceptions to its general enactments, always baffled the efforts of her legislators.

Thus it is with the classification of the animal, the vegetable, and the mineral, kingdoms. Ingenious distinctions have been laid down for the discrimination of one family from another, but in spite of our science in defining divisions, anomalous subjects continually step in, and show that our system is unfounded in nature. Indeed, how can we hope to class, with precision, the component parts of these several kingdoms, when human science is inadequate to point out a clear line of distinction between the three kingdoms themselves.

Nevertheless, the present state of the science is truly inestimable to us. Men, the most learned and industrious, have contributed to its present perfection; and talents, the most eminent, have been devoted to its extension. Much has been attained, and still much we may hope to discover, notwithstanding the whole depth of the subject is alone fathomable by him who framed its laws.

This magnificent plant has frequently been imported from Constantinople, and received its trivial appellation, *Susiana*, from a western district of Persia, which was known to the ancients by that name. Susa, now called Caster, was the capital of that district, and the term, in the language of the country, signifies Lilies. Chalcedon, a city of Asia, opposite Constantinople.

It requires to be planted in a warm, dry, loamy, or gravelly soil, or it will rarely be found to blossom freely. To preserve the plant in health, it seems desirable to transplant it once in four or five years, as parts of the old roots undergo decay, and become prejudicial to the growth of the young ones.

Hort. Kew. 2, v. 1, 117.

LILIUM BULBI-FERUM.

BULB-BEARING LILY.

Class.
HEXANDRIA.

Order.
MONOGYNIA.

Natural Order.
LILIACEÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
Italy.	3 feet.	May, June.	Perennial.	in 1596.

No. 31.

LEIRION is the Greek name of the Lily; and from it, through the Latin, our term Lilium has been derived.

Much has been said on the analogy between bulbs and buds, and their affinity is more perfect than many of our readers probably imagine.

This species of Lily presents a pretty example of the bulb produced in the usual situation of the bud, and the affinity thereby becomes more evident. We shall treat this subject more at length in a future number.

The culture of this plant is extremely easy. The old roots increase very little, but the bulbs taken from the stem, in August, and then deposited in the soil, will produce one leaf in the following spring: in the second year, a small bulb-bearing stem, about fifteen inches high; but not usually a flower: in the third year, a stronger stem, bearing bulbs, and terminated by one of its beautiful blossoms; and a handsome display of flowers may be expected in the following summer. Transplanting should be effected in autumn; and shade or exposure will be equally suitable.

Hort. Kew. 2, v. 2, 241.

SPIRÆA LEVIGATA.

SMOOTH SPIRÆA.

Class.
ICOSANDRIA.

Order.
DI-PENTAGYNIA.

Natural Order.
ROSACEÆ.

Native of Siberia.	Height. 4 feet.	Flowers in Apr. June.	Duration. Perennial.	Cultivated in 1774.
-----------------------	--------------------	--------------------------	-------------------------	------------------------

No. 32.

The term *Spiræa* has been deduced from the Greek *SPEIRA*, a rope; and is applied to this genus in consequence of the flexibility of the branches in some of its species. *Lævigata*, from the Latin *lævis*, smooth.

There are several species of *Spiræa* well known in our gardens, as the *Spiræa salicifolia*, usually called the *Spiræa frutex*; and the *Spiræa hypericifolia*, sometimes known by the name of *Hypericum frutex*. Our present subject is not quite so frequently met with as either of these, though not less interesting, nor less entitled to a place in every shrubbery. Its summer foliage is generally retained in a dry state, during winter, and forms a nice shelter to the young shoots, which are usually rather succulent, and seem to need some such protection against the inclemencies of winter.

It may occasionally be propagated by suckers, though not so freely as are some others of the same family. Or it may readily be raised by layers, which should be confined under the soil in autumn, and they will be sufficiently rooted for separation in twelve months.

Hort. Kew. 2, v. 3, 254.

SPINOSA LEVIGATA.

SMOOTH SPINOSA.

Order.
DIAPYCNIA.

Class.
LEGUMINOSAE.

Family.
SPINOSA.

Native of	Height	Flowers in	Quantity	Coloured
Siberia	4 feet	the June	terminal	in 1771


W. 52.

The term *Spinosa* has been deduced from the Greek *spinē*, a spin; and is applied to this genus in consequence of the flexibility of the branches in some of its species. *Levigata*, from the Latin *laevigatus*, smooth.

There are several species of *Spinosa* well known in our gardens, as the *Spinosa rubra*, usually called the *Spinosa frutes*; and the *Spinosa hypoleuca*, sometimes known by the name of *Hypoleuca frutes*. Our present subject is not quite so frequently met with as either of these, though not less interesting, nor less entitled to a place in every shrubbery. Its summer foliage is generally retained in a dry state, during winter, and forms a nice shelter to the young shoots, which are usually rather succulent, and seem to need some such protection against the inclemencies of winter.

It may occasionally be propagated by cuttings, though not so freely as are some others of the same family. Or it may readily be raised by layers, which should be covered under the soil in autumn, and they will be sufficiently rooted for separation in twelve months.

Bot. Rev. 2, 1, 254.





Kalmia latifolia.



Linaria purpurea.



Oenothera pumila.

$\frac{1}{2}$



Valeriana montana.

$\frac{1}{2}$

KAL'MIA LATIFO'LIA.

BROAD-LEAVED KALMIA.

Class.

DECANDRIA.

Order.

MONOGYNIA.

Natural Order.

RHODORACEÆ.

Native of N. America.	Height. 4 feet.	Flowers in May, July.	Duration. Perennial.	Introduced in 1734.
--------------------------	--------------------	--------------------------	-------------------------	------------------------

No. 33.

Peter Kalm was an eminent botanist of Finland, and professor of the science at Abo. He imbibed a love of nature among the flowery rocks of Sweden; the wilds also of North America, and icy forests of Russia, were alike explored by him. He published his travels through North America in 1753, which were translated into English by Mr. Forster in 1771; and Linneus, it may be conceived, paid no more than due respect to merit, in bestowing his name on this splendid family of plants. Latifolia, from the Latin latus, broad; and folium, a leaf.

One sentiment, usually excited by the presence of this beautiful North American shrub, is that of regret, arising out of the difficulty of keeping it in perfect health. The principal requisites towards this appear to be purity of air, moisture and shade during our summer months, and a proper imitation of its native soil.

Abercrombie says, 'most of the exotic shrubs brought from America, were originally found growing on tracts of ground resembling our beds of peat, and the luxuriance of these vegetables may partly be ascribed to the excessive moisture which is

peculiar to some parts of America. One great object is to imitate the American peat. This is a composition of the branches, twigs, leaves, and the roots of trees; with small plants, grass, and weeds; by having lain immemorially in water, the whole is formed into a soft mass; and when the materials are completely decayed, and blended so as to be homogeneous in appearance, the compound is the finest vegetable mould: where this description of peat cannot be obtained, recourse must be had to the best that can be procured from marshes, bogs, or heathy commons, which must be well turned and sweetened, and mixed with sand and rotten leaves.'

Of this soil, distinct beds should be formed, about three feet deep, for this class of American plants, with a shade of shrubby or lofty trees on the south side; and if the beds could be so placed as to admit an occasional flow of water being turned through a stratum of pebbles, laid three or four feet beneath the surface, little doubt would exist of success.

It is preferable to transplant in spring or autumn, but it may be effected at almost any period, if the earth be not too much disturbed from the fibrous roots, which are so small and numerous as to retain the soil in one compact mass. But little difficulty would exist in the cultivation of the *Kalmia*, *Rhododendron*, and other similar plants, if a sufficiency of moisture could be conveniently afforded them during our summer months. In most situations they require copious waterings in very warm weather.

Our drawing is from the pale-flowered *Kalmia latifolia*, and we intend resuming the subject, with further directions, under another variety of it.

Hort. Kew. 2, v. 3, 47.

LINARIA PURPUREA.

PURPLE FLAX-WEED.

Class.
DIDYNAMIA.

Order.
ANGIOSPERMIA.

Natural Order.
SCROPHULARINÆ.

Native of S. Europe.	Height. 4 feet.	Flowers in June, Sep.	Duration. Perennial.	Cultivated in 1648.
-------------------------	--------------------	--------------------------	-------------------------	------------------------

No. 34.

The term *Linaria*, appears to have had its origin from *linum*, *flax* : just as *Flax-weed* and *Toad-flax* are deduced from the English word, on account of the similarity of the plants.

This species is frequently known as the *Antirrhinum purpureum*, in which genus it was placed by *Linneus* : modern botanists have, however, divided the *Antirrhinums* into two genera, and this plant now stands with the appellation by which it was well known to *Gerard*, *Parkinson*, *Bauhin*, and others. *Purpurea*, from the Latin, purple.

Its erect and elegant growth renders it well suited to contrast with more diffuse subjects of its own stature; and the simplicity of its culture will qualify its deficiency of that splendour which may attach to some of its more fastidious neighbours.

It produces seed freely, and from these may be readily propagated. If sown in the autumn, upon a tolerably dry soil, they will not fail to come up, and produce much stronger plants than those sown in spring; and when once established it may be expected to increase, by seed, spontaneously, though in very wet soils it will sometimes fail.

Hort. Kew. 2, v. 4, 12.

ÆNOTHE'RA PU'MILA.

DWARF ÆNOTHERA.

Class.
OCTANDRIA.

Order.
MONOGYNIA.

Natural Order.
ONAGRARIÆ.

Native of N. America.	Height. 1 foot.	Flowers in May, Aug.	Duration. Perennial.	Cultivated in 1757.
--------------------------	--------------------	-------------------------	-------------------------	------------------------

No. 35.

From two Greek words OINOS, and THERA, the first signifying wine, and the second a pursuit, in consequence of the dried roots acquiring the flavour of wine. Pumila, from the Latin, little.

Ænothera has, by different authors, been accented on the antepenultimate, and on the penultimate, syllable: the latter agrees best with the derivation.

Though this is the smallest Ænothera that we are acquainted with, it forms a pretty ornament amongst rock-work, and plants of like size; and we find it possess this advantage, that whilst blossoming through the hottest months, it rarely suffers by drought.

It is easily increased by parting the roots in spring or autumn; but it will produce much finer plants, and flower better, when raised from seeds. Miller directs that they be sown in autumn, in pots, placed under a hotbed frame until the spring, when the plants will appear: when fit to remove, plant a few in small pots, to be sheltered under a common frame in winter; others may be set in a sheltered border, and the following summer they will produce flowers and seeds in plenty.

Hort. Kew. 2, v. 2, 343.

GEOTHEKA PIPERITA.

DWARF GEOTHEKA.

Geotheca
Piperita

Geotheca
Piperita

Geotheca
Piperita

Native of	Height	Flowers in	Flowers in	Collected
N. America.	1 foot	May, Aug.	June, Sept.	in 1857.

No. 22.

From two Greek words *geos*, and *thera*, the first signifying wine, and the second a parent, in consequence of the dried roots acquiring the flavor of wine. *Piperita*, from the Latin, little.

Geotheca has, by different authors, been accounted on the antipathetic, and on the penicillate, syllable: the latter agrees best with the derivation.

Though this is the smallest *Geotheca* that we are acquainted with, it forms a pretty ornament amongst rock-work, and plants of like size; and we find it possesses this advantage, that whilst blossoming through the hottest months, it rarely suffers by drought.

It is easily increased by putting the roots in spring or autumn; but it will produce much finer plants, and flower better, when raised from seeds. Miller directs that they be sown in autumn, in pots, placed under a hotbed frame until the spring, when the plants will appear: when fit to remove, plant a few in small pots, to be sheltered under a common frame in winter; others may be set in a sheltered border, and the following summer they will produce flowers and seeds in plenty.

Herb. Kew. 2, v. 2, 242.

VALERIA'NA MONTA'NA.

MOUNTAIN VALERIAN.

Class.
TRIANDRIA.

Order.
MONOGYNIA.

Natural Order.
DIPSACEÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
Switzerland	1 foot.	May, June.	Perennial.	in 1739.

No. 36.

Of the origin of the term Valeriana, we have nothing certain on record. Some persons have supposed it to have originated in the name of an eminent physician, Valerius, who is said to have first used one of the species of it in medicine; whilst others think the term may have come from the Latin valere, to be well. Montana, from the Latin mons, a mount.

We are not aware that this species has been employed in medicine, but the Valeriana officinalis is not only used against particular disorders; but, according to Gerard, was in his day employed as a pot herb by the inhabitants of the north. He quotes a lame couplet in its praise, and says, 'some woman poet or other hath made these verses.' Could this venerable herbalist be introduced to some of our 'woman poetry' of the nineteenth century, we think he would not, so unceremoniously, cast a slur on the productions of the fair sex.

It flourishes in a light dry soil, and is readily increased by a separation of its roots. Though increase be not wanted, it will still be desirable to divide and transplant it occasionally.

Hort. Kew. 2, v. 1, 74.

VALERIANA MONTANA

MOUNTAIN VALERIAN.

Gray.
MONTANA

Gray.
MONTANA

Gray.
MONTANA

Height of Stems 1 foot.	Flowers in May, June.	Collected in 1878.
----------------------------	--------------------------	-----------------------


No. 25

Of the origin of the term Valerian, we have no-
thing certain on record. Some persons have sug-
posed it to have originated in the name of an ancient
physician Valerius, who is said to have first used
one of the species of it in medicine; whilst others
think the term may have come from the Latin va-
lere, to be well. Montana, from the Latin mons, a
mountain.

We are not aware that this species has been em-
ployed in medicine, but the Valerian officinalis is
not only used against hysterical disorders; but ac-
cording to Gerard, was in his day employed as a pre-
servative by the inhabitants of the north. He quotes a
Latin receipt in his process, and says, 'some women
put on other herbs made thus warm.' Could this
valerian be the one introduced in some of our
'woman's penny' of the nineteenth century, we think
it would not, so unceremoniously, cast a slur on the
production of the fair sex.

It sometimes is a light dry soil, and is readily in-
creased by a repetition of its roots. Though increase
be not wanted, it will still be desirable to divide and
replant it occasionally.

How to use it, p. 14.





Colutea frutescens

2/5



Gnaphalium arenarium

3/5



Hesperis matronalis

2/5



Senecio elegans

3/5

COLUTEA FRUTESCENS.

SCARLET BLADDER SENNA.

Class.
DIADELPHIA.

Order.
DECANDRIA.

Natural Order.
LEGUMINOSÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
C. G. Hope.	4 feet.	June, Aug.	Perennial.	in 1683.

No. 37.

The term, Colutea, has been introduced from the Greek name of a plant in Theophrastus, but no further knowledge appears to exist of its origin or application. It probably comes from ΚΟΙΛΟΣ, a spacious cavity, in allusion to its distended seed vessel. Frutescens, from the Latin frutex, a shrub. The common term Bladder Senna, has been given to it from its bladder-like pods, and the general resemblance of its foliage to that of the Official Senna. It is also said to possess a portion of the cathartic qualities of that plant.

Colutea is one of the unfortunate genera, among many others, that has come, perfected as it were, from the hand of the great father of our artificial system, to suffer amputation and distribution in after ages. The Colutea frutescens is now found in the modern catalogues as Sutherlandia frutescens, in consequence of this name being adopted in the Hortus Kewensis, after Mr. Robert Brown. To the very eminent abilities of the author of the Prodromus Novæ Angliæ, every one must be ready to pay the tribute of praise; but, perhaps, that very acuteness of observation, and depth of botanical

science for which he is celebrated, may render him more nice in his generic divisions than the plain botanist may admit to be necessary, or, indeed, than is useful to the practical man; for certain it is, that the extension of our botanic vocabulary, and perplexing increase of synonyms, form a considerable drawback on any advantage that may accrue from nicer distinctions. Both Sir J. E. Smith, and Dr. Sims are of opinion, that the genera *Sutherlandia* and *Swainsonia*, are too nearly allied to *Colutea* to admit separation.

Its beautiful scarlet flowers, contrasted with its silvery foliage, render this a peculiarly ornamental little shrubby plant; and after its brilliant blossoms are faded, its large inflated pods still excite an interest in its welfare. It is rather tender, and is frequently kept in the greenhouse, where it never blossoms so luxuriantly as in the open air.

It should be planted against a wall, in a warm sheltered situation, in light dry soil; and in frosty weather be protected by litter and matting. Or if kept in a pot, in the house, it should be placed near to the window; and it will always be benefited by being fully exposed in mild weather.

Treated as a hardy plant, it does not continue more than three or four years: this, however, is not important, since it is easily raised from seeds, which may be sown early in the spring, in pots, placed in a cool cucumber frame, and the plants should be hardened to the open air as early as possible. If thus forwarded, they will sometimes blossom in the latter part of the summer; or, if not at that time, early in the following season.

Hort. Kew. 2, v. 4, 327.

GNAPHA'LIIUM ARENA'RIIUM.

SAND EVERLASTING.

Class.

SYNGENESIA.

Order.

POLYGAMIA SUPERFLUA.

Natural Order.

CORYMBIFEREÆ.

Native of Europe.	Height. 9 inches.	Flowers in June, July.	Duration. Perennial.	Cultivated in 1739.
----------------------	----------------------	---------------------------	-------------------------	------------------------

No. 38.

The name comes from the Greek GNAPHALON, signifying down or wool, such as is produced by dressing or shearing cloth, in allusion to the woolliness of the foliage. Arenarium, from the Latin arena, sand, indicative of the soil which it prefers.

The presence of the flowers of this Gnaphalium, in many parts of Europe, gives rise to feelings of melancholy and sadness, which Englishmen have never been taught to connect with them. The contrary is rather the case, for they are introduced among us in bouquets for festivities, and embellishments for head dresses. They are sometimes artificially tinged with a diversity of hues; and from their continued beauty, in a dry state, they have obtained the title Everlasting.

Some of our readers may not be informed of the prevalent practice, amongst the inhabitants of the continent, in using these flowers to decorate the monuments and graves of their departed relatives and friends. The French are particularly partial to these flowers, and designate them as we do, IMMORTELLE. It has been observed by Mr. Phillips, that 'since the hill of Pere la Chaise has been

converted into a cemetery for the city of Paris, the demand for these flowers in the French capital has been so considerable, as not only to employ many hands in the cultivation of them, but numerous families are regularly occupied, and entirely supported by forming these flowers into garlands and crosses, which are offered for sale by the cottagers near the entrance of this celebrated burial ground; and but few persons can visit the romantic and hallowed spot, without having some name called to their remembrance, which draws from them this slight token of remembrance; for here we find a mingled mass of monuments, recalling to our recollection the sweet lines of the poet, the ready wit of the critic, the piety of the priest, the heroic deeds of the soldier, the bravery of the sailor, the labours of the naturalist, the beauties of the artist, and the loves of Abelard and Heloise; here we meet fond parents with wreaths of IMMORTELLS to drop on the sod of their blighted hopes, and affectionate children placing crosses of everlasting flowers on the head of their parents' graves.'

The cultivation of this plant is particularly easy, where the soil is light and dry; but in damp situations, it frequently dies, under the influence of the moisture of our climate during the first three months of the year. It should be divided in spring or autumn, and if the soil be stiff, mix with it an ample portion of drift sand; this will generally preserve it in health; but should it not do so, plant it on little hillocks of light soil, which may be raised six or eight inches high, as a further defence against damp; and half cover the soil with small pebbles.

Hort. Kew. 2, v. 5, 14.

HESPERIS MATRONALIS. plena alba.

DOUBLE WHITE ROCKET.

Class.
TETRADYNAMIA.

Order.
SILIKUOSA.

Natural Order.
CRUCIFERÆ.

Native of Italy.	Height. 18 inches.	Flowers in June, July.	Duration. Perennial.	Cultivated in 1597.
---------------------	-----------------------	---------------------------	-------------------------	------------------------

No. 39.

Hesperis, from the Greek HESPEROS, evening; in consequence of the fragrance of some of this tribe at that time. We are told that the ladies of Germany cultivate these flowers in pots, to secure the evening perfume of them in their apartments, whence they have been called dames' violets; and this allusion has been Latinized into matronalis. The name of Rocket seems to have come to us through the Latin eruca, signifying canker worm, as well as the name of this plant, and as it frequently dies with the unskilful, without a perceptible cause, the term may be as applicable as many others in our scientific nomenclature.

Many and contradictory directions have been given for the cultivation of this plant, evidently by persons not at all practically acquainted with its habits. The best information we have ever seen printed, is contained in Rees's Cyclopædia. The grand secret consists in cutting down the flower stems of a plant or two, before they are much exhausted by the bloom; thus a good stock of offsets will be produced. A cool and rather moist situation is always to be preferred for the rocket.

Hort. Kew. 2, v. 4, 122.

SENE'CIO E'LEGANS.

PURPLE GROUNDSEL.

Class.

SYNGENESIA.

Order.

POLYGAMIA SUPERFLUA.

Natural Order.

CORYMBIFEREÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
C. G. Hope.	2 feet.	June, Oct.	Annual.	in 1700.

No. 40.

Senecio, from the Latin senex, an old man; a name which has been given to it from the fancied resemblance of its seed down to the grey or silvery head of age. Elegans, elegant. The seed of this plant is usually sold under the name of Jacobæa, which term has been handed down from some old botanists, who called it Flos Sancti Jacobi, or the flower of Saint James.

The Senecio elegans may be raised from seeds, sown with other hardy annuals in the spring, and requires no peculiar care.

The double one, which is merely a variety of the same, is an extremely beautiful plant; and although it does not come within our limits as a hardy one, we may be pardoned the union of it with its synonymous species. It is usually cultivated in pots, as an ornament for the house, yet few plants are better adapted for giving assistance to the brilliancy of the flower border, where, during the summer months, it will grow with greater luxuriance and beauty than in any other situation.

It is a tender plant, and where there is not a greenhouse protection, it may be kept during the

winter, in a dry airy room, protected from frost ; and should be very sparingly supplied with water, or its succulent stems will be liable to decay. In the beginning of March, propagation may be commenced by cuttings, and if one plant only has been preserved through the winter, an abundance of others may be raised from it. Two joints are sufficient to constitute a cutting, and it should be taken off close beneath the lower one. It is usual to slit it a quarter of an inch upwards from the bottom, through the joint, which certainly in some plants facilitates their rooting, and can injure none. These should be planted in pots of light rich earth, two or three in each, then watered immediately ; and bell glasses, or in the absence of these, tumblers or goblets, be turned over them. If in this state it be convenient to place the pots in a warm cucumber frame, the cuttings will strike root more quickly ; but if not, they may be placed before the window of a warm room, having a southern aspect, where they will generally succeed very well. The glasses should, occasionally, be taken from them for a few minutes, and wiped, but not be altogether removed till the cuttings have struck root. When they begin to grow freely, they may be transplanted into separate pots, be gradually hardened to the open air, and kept in readiness to turn into the borders about the middle of May, where they will prove a conspicuous ornament all the summer. The white variety of the Double Groundsel is far less common than the purple ; and though not so gay an ornament, is a very desirable one, either for pot culture or to beautify the parterre.





Oenothera macrocarpa.

70



Pentstemon pubescens.

71



Kalmia glauca.

E. D. Smith del.

72



Achillea speciosa.

73

S. Watts sculp.

ŒNOTHERA MACROCARPA.

LARGE-FRUITED EVENING PRIMROSE.

Class.
OCTANDRIA.

Order.
MONOGYNIA.

Natural Order.
ONAGRARIÆ.

Native of	Height.	Flowers in	Duration.	Introduced
N. America.	18 inches.	June, Aug.	Perennial.	in 1811.

No. 41.

Œnothera, from the Greek, explained under Œnothera pumila, No. 35. Macrocarpa is also of Greek extraction, signifying large-fruited.

Nearly the whole of the genus Œnothera, or Evening Primrose, a name by which some of the species are so familiar to us, are extremely ornamental plants. They have obtained the latter appellation from the circumstance of their flowers expanding in the evening; though this peculiarity is not equally applicable to all.

The species Macrocarpa has been considered synonymous with Missourensis.

It is easily propagated by cuttings, which may be planted in pots, and placed in a hotbed, or under a hand glass, till they have struck root; they will there be convenient for removing into warm parts of the flower border, or to place in a cold frame for more effectual winter protection.

It should be planted in a situation favourable to its trailing on the ground, unless considerable attention can be paid to the support of its succulent stems; and even then it will be less pleasing than when naturally spreading on the earth.

CHOTHEVA MAYOCAPPA

LEAF-THEFT EVENING PRIMROSE

1887
MAYOCAPPA

1887
MAYOCAPPA

Mayocappa
Mayocappa

Native of	Height	Flowers in	Lat. - Long.
18 inches	10 inches	10 inches	10 inches

No. 11

Chotherva, from the Greek, epithet under
Chotherva family, No. 11. Chotherva is the
Greek epithet, signifying large-fruited.
Nearly the whole of the plant, as
the young primrose, a name for which some of the
specimens are so similar to the extremely small
mossy plants. They have obtained the latter ap-
plication from the circumstance of their flowers ex-
posed in the evening; though this peculiarity
is not equally applicable to all.

The species Chotherva has been considered
as a synonym with Chotherva.
It is easily propagated by cuttings, which may
be placed in pots, and placed in a hotbed, or under
a glass pane, all they have to do; they will
not be convenient for removing into cold pots at
the winter border, or in place in a cold house for
more effectual winter protection.

It should be placed in a situation exposed to
the sun on the ground, where considerable frost
may be laid to the surface of the ground; and
even then it will be less injured than when in
the earth.

For May 1 1887

PENTSTEMON PUBESCENS.

HAIRY PENTSTEMON.

Class.
DIDYNAMIA.

Order.
ANGIOSPERMIA.

Natural Order.
BIGNONIACEÆ.

Native of N. America	Height. 18 inches.	Flowers in July, Sept.	Duration. Perennial.	Cultivated in 1758.
-------------------------	-----------------------	---------------------------	-------------------------	------------------------

No. 42.

From two Greek words, united to signify a fifth stamen; which name has been chosen to distinguish this genus from others of the same class, though the additional stamen is always more or less imperfect. Pubescens, from the Latin, downy.

It is frequently known as the Chelone pentstemon, and this, some of the best botanists still consider its legitimate title; but the use of the new name, amongst those who never even pretend to judge of its propriety, has assisted in establishing the division of it from Chelone. It was first adopted merely on account of a trifling variation in the rudiment of the additional filament, already alluded to.

It requires but little care in cultivation, and may be propagated either by seeds, cuttings, or separation at the root. Seeds should be sown in the spring, and the plants, which require no other care than that of being kept free from weeds, will blossom in the following summer. If cuttings be taken as early in the spring as the plants will admit, they will flower in the following autumn; and as the old roots are liable to decay, frequent propagation will be requisite.

Hort. Kew. 2, v. 4, 8.

PLANTATION HISTORY

PLANTATION HISTORY

PLANTATION HISTORY

PLANTATION HISTORY

PLANTATION HISTORY	PLANTATION HISTORY	PLANTATION HISTORY	PLANTATION HISTORY	PLANTATION HISTORY
PLANTATION HISTORY	PLANTATION HISTORY	PLANTATION HISTORY	PLANTATION HISTORY	PLANTATION HISTORY

PLANTATION HISTORY

From two Greek words, *plantatio* and *historia*, the name *plantation* has been chosen to distinguish the genus from other of the same class, though the additional element is always more or less important. *Plantatio*, from the Latin, *planta*, is a frequently known name in the botanical literature, and that, some of the best authorities will consider its botanical title; but the use of the word *plantation* is more common than with other even proposed to judge of its propriety, has resulted in establishing the division of it from *Chlorophyta*. It was first adopted nearly in its present of a striking variation in the treatment of the additional element, already alluded to. It requires but little care in cultivation, and may be propagated either by seeds, cuttings or layers, and of the root. Good should be seen in the spring, and the plants, which require no other care than that of being kept from weeds, will blossom in the following summer. It is called *plantatio* in the Latin as the spring as the plants will bloom, they will flower in the following autumn, and as the old texts are liable to decay, frequent propagation will be required. *Plantatio* is the Latin name of the *plantatio*.

PLANTATION HISTORY

KALMIA GLAU'CA.

GLAU'COUS KALMIA.

Class.

DECANDRIA.

Order.

MONOGYNIA.

Natural Order.

RHODORACEÆ.

Native of N. America.	Height. 3 feet.	Flowers in April, May.	Duration. Perennial.	Cultivated in 1767.
--------------------------	--------------------	---------------------------	-------------------------	------------------------

No. 43.

Kalmia, after Peter Kalm, noticed at No. 33, where a return to the same subject was anticipated under another variety of Kalmia latifolia. The Kalmia glauca is now given, there being but little difference in the nature of the several species. In America it is called the spoon tree, in consequence of the natives making use of the wood, which is close and hard, for the purpose of forming various articles of household utility.

Miss Edgeworth, in her tale of "To-morrow," ingeniously alludes to the poisonous properties of the leaves of this splendid plant. It forms one of her auxiliaries in the delineation of the evils of procrastination, which she has given with a force and feeling not to be surpassed. The incident is founded on the communication of Dr. Barton, in the American Transactions; who states that in Philadelphia, in the winter of 1790, an official proclamation was issued, warning all persons from eating pheasants. This, as Dr. B. states, was done on a well grounded opinion, that several persons had died from the use of these birds, which had fed on the leaves of the Kalmia. Opposed to this opinion is that of

Mr. Wilson, the ornithologist, who says that he has found the crops of these birds distended almost entirely with the buds of the *Kalmia*, but that he has eaten freely of the flesh of such birds without experiencing any ill consequence.

Its noxious character has been strongly exhibited on the rattle snake. A few drops of a tincture prepared from the leaves, having been poured on this reptile, it died shortly afterwards: and the deleterious effects of the leaves on the human system, are such as should excite a strict guard against their being eaten by children.

Kalm, himself, states that both sheep and calves have died from eating them; and that cows and horses had suffered greatly from the same cause. On the other hand, when the ground is covered with snow, stags browse them as their common food, not only without injury to themselves, but also without any ill effects having been known to arise from the use of the venison.

In the cultivation of plants generally, but more particularly those of difficult growth, attention should always be given to the peculiar situation of them in their native soil. Professor Kalm has chiefly noticed the broad-leaved species, and found it succeed best on the sides of hills, especially on the north side; and says, on meeting with a steep place near a brook, or on the side of a hill, towards a marsh, you are sure to find it. This alone speaks pages to the attentive cultivator, particularly when he is further told, that vegetable earth or peat, constitutes its native soil; and that in a climate very similar to our own.

Hort. Kew. 2, v. 3, 47.

ACHILLE'A SPECIO'SA.

SPEAR-LEAVED MILFOIL.

Class.
SYNGENESIA.

Order.
POLYGAMIA SUPERFLUA.

Natural Order.
CORYMBIFERÆ.

Native of	Height.	Flowers in	Duration.	Introduced
Unknown.	3 feet.	July, Aug.	Perennial.	in 1804.

No. 44.

The name Achillea, is deduced from Achilles, the famous Grecian hero, whose strength and prowess forms so conspicuous a character in the Iliad. Clad in armour, impenetrable and resplendent, Homer compares him with the sun;---

“Not brighter, Phœbus in the ethereal way,
Flames from his chariot and restores the day.”

Speciosa, from the Latin, signifying handsome. Many of the plants now coming under the denomination of Milfoil, do not possess the character which this name seems to have been first intended to indicate; the Latin millefolium, of which it is a corruption, being compounded of mille, a thousand; and folium, a leaf; from the very numerous pinnæ, or leaflets, on some of these plants.

The flowers of this Achillea are, probably, not so showy and attractive as the florist may be led to expect from its appellation of Speciosa. The plant, however, is of handsome upright growth, and its serrated foliage particularly neat. It is of easy culture, and admits a division of its roots at the usual season. These generally increase sufficiently to afford an ample supply.

Loudon's Ency. of Plants, sp. 12352.

ACHILLEA STECIOSA

and the name of the plant is Achillea steciosa.

It is a perennial herb, with a thick, woody root.

The leaves are narrow, lanceolate, and serrated.

The flowers are small, white, and numerous.

It is found in dry, open places, and is common in the mountains.

The name Achillea is derived from the name of the hero Achilles.

The name steciosa is derived from the Greek word stekios, meaning thick or woody.

The plant is used in medicine for its astringent and tonic properties.

It is also used in the preparation of various wines and liquors.

The plant is very hardy and can withstand severe frosts.

It is a very common plant in the mountains of the Alps.

The name Achillea is also given to other plants of the same genus.

The name steciosa is also given to other plants of the same genus.

The plant is very common in the mountains of the Alps.

The name Achillea is also given to other plants of the same genus.

The name steciosa is also given to other plants of the same genus.

The plant is very common in the mountains of the Alps.

The name Achillea is also given to other plants of the same genus.

The name steciosa is also given to other plants of the same genus.

The plant is very common in the mountains of the Alps.

The name Achillea is also given to other plants of the same genus.

The name steciosa is also given to other plants of the same genus.

The plant is very common in the mountains of the Alps.

The name Achillea is also given to other plants of the same genus.

The name steciosa is also given to other plants of the same genus.

The plant is very common in the mountains of the Alps.

The name Achillea is also given to other plants of the same genus.

The name steciosa is also given to other plants of the same genus.

The plant is very common in the mountains of the Alps.

The name Achillea is also given to other plants of the same genus.

The name steciosa is also given to other plants of the same genus.

The plant is very common in the mountains of the Alps.

The name Achillea is also given to other plants of the same genus.

The name steciosa is also given to other plants of the same genus.

The plant is very common in the mountains of the Alps.


The name Achillea is also given to other plants of the same genus.

The name steciosa is also given to other plants of the same genus.

The plant is very common in the mountains of the Alps.

The name Achillea is also given to other plants of the same genus.

The name steciosa is also given to other plants of the same genus.





Verbascum Phoeniceum.

2/3



Coreopsis lanceolata.

1/2



Ledum palustre.



Campanula patula.

VERBASCUM PHŒNICEUM.

PURPLE-FLOWERED MULLEIN.

Class.
PENTANDRIA.

Order.
MONOGYNIA.

Natural Order.
SOLANÆÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
S. Europe.	4 feet.	May, Aug.	Perennial.	in 1596.

No. 45.

Verbascum seems to be of very uncertain derivation, and the term having been used by the oldest writers, its origin is likely to remain in obscurity. A supposition has been ventured that it is a corruption of barbascum, from barba, bearded, on account of the woolliness of some of the species. Phœniceum, from the Latin, signifying purple-coloured.

Of the origin or application of our English term Mullein, we find no traces, otherwise than as the name of a plant. Some of these being soft and woolly, as before observed, the word very probably, has been adopted from the Latin mollis, as indicative of that quality.

With our drawing of this plant we have given a representation of a section of its stem. On cutting the stem through transversely, the arrangement of its sap vessels are as perfectly exhibited as in vegetables of a more ligneous formation; and it is pretty certain, that its functions of vegetable life are performed by ascending and descending fluids upon the selfsame principles.

On the circulation of vegetable fluids, various theories have, at different periods, been promulgated

by ingenious physiologists. Their theoretical speculations, however, have been pursued with much abstruseness and uncertainty, and it was left for the present age to exhibit, with somewhat like precision, the laboratory of nature in the vegetable kingdom. Mr. Knight, the president of the Horticultural Society, from thirty years' intense application to this subject, has determined, by ingenious and satisfactory experiments, many of the phenomena of vegetation, particularly such as are connected with the circulation of the sap, and the perspiration of plants.

These subjects, perhaps, some of our readers may not have had occasion to examine. As they cannot fail to prove of peculiar interest to many inquiring minds, we shall have pleasure, as opportunity offers, in giving the opinions of eminent botanists, and the result of various experiments connected with this department of vegetable physiology.

This species of *verbascum* is particularly handsome, and we have observed that the flowers of a plant which has remained in its present situation four or five years, are now produced of a darker hue than when it was first planted. This possibly may have arisen from the presence of a portion of peat soil, which, from an alkaline quality that exists in it, is sometimes found to change pinks into purples; perhaps from the neutralization of acids, which produce effects directly opposite.

It never has perfected seeds with us, nor does it increase very much at the root, but it may be propagated by cuttings of the young flowering stems, which strike root readily under bell glasses. It does not require any peculiar soil or situation.

Hort. Kew. 2, v. 1, 385.

COREOPSIS LANCEOLATA.

SPEAR-LEAVED COREOPSIS.

Class.
SYNGENESIA.

Order.
POLYGAMIA SUPERFLUA.

Natural Order.
CORYMBIFERÆ.

Native of Carolina.	Height. 3 feet.	Flowers in July, Oct.	Duration. Perennial.	Introduced in 1724.
------------------------	--------------------	--------------------------	-------------------------	------------------------

No. 46.

The plants of this genus are sometimes called the tick-seed sun flower, in allusion to the shape of their seed; which circumstance is regarded in the systematic appellation, coreopsis, noticed under No. 13. Lanceolata, from the form of the leaves, which approaches that of a lance or spear.

There are very few plants which exhibit a more rich profusion of golden flowers, and also produce them as long in succession as the Coreopsis lanceolata. From its medium height it is well suited either to the borders allotted to the growth of herbaceous subjects, or for introduction into the mingled plantation of low flowering shrubs, where it will form a conspicuous ornament.

It may be increased by dividing the roots in spring or autumn, but preference should be given to the former season; or cuttings may be taken at any time during the summer months, provided it be sufficiently early to admit of their making strong root before the commencement of cold weather. Not less than two months should be allowed for this purpose, even with the assistance of bell or hand glasses.

Hort. Kew. 2, v. 5, 135.

LEDUM PALUSTRE.

MARSH LEDUM.

Class.
DECANDRIA.

Order.
MONOGYNIA.

Natural Order.
RHODORACEÆ.

Native of Europe.	Height. 3 feet.	Flowers in April, May.	Duration. Perennial.	Introduced in 1762.
----------------------	--------------------	---------------------------	-------------------------	------------------------

No. 47.

The name of this genus has been adopted from LEDON, which the Greeks applied to a species of cistus that produced their LEDONON, our labdanum, a resinous substance, sometimes employed in plasters; and also in fumigations, for its perfume. The term has been improperly referred to lædendo as its original. Palustre, from the Latin, marshy, in allusion to its native situation.

This plant, when bruised, has a rather pleasant aromatic scent, similar to that of fresh gathered hops; and Linneus tells us, that the inhabitants of some parts of Sweden, make use of the leaves in their beer, which produce an agreeable flavour, but an intoxicating quality.

The Laplanders are said to strew the branches amongst their grain to drive away mice.

A little variety occurs in the colour of its flowers. Some are of a clear white; whilst others are delicately tinged with a pale pink.

The Ledums come under the denomination of peat plants; and grow best in that soil. This species succeeds best in a shady situation; and in the summer, should be supplied with copious waterings.

Hort. Kew. 2, v. 3, 48.

LEOPOLD PATENT

NEW METHOD OF CULTIVATING

Order
Address

Order
Address

Patent Office
London

Height of Plant	Height of Plant	Height of Plant	Height of Plant
1 foot	2 feet	3 feet	4 feet

No. 1

The name of this grape has been adopted from
arrows, which the Greeks applied to a species of
vine that produced their arrows or, our Indians,
a resinous substance, sometimes employed in pain-
tens, and also in fumigations, for its perfume. The
term has been improperly referred to Indians as its
original. Perhaps, from the Latin, *paris*, in allu-
sion to its native situation.

This plant, when trained, has a rather pleasant
aromatic scent, similar to that of fresh gathered
leaves; and I have seen that the inhabitants of
some parts of Sweden, make tea of the leaves in
their beer, which produces an agreeable flavour, but
an intoxicating quality.

The Laplanders are said to brew the branches
amongst their grain to drive away vermin.

A little variety occurs in the colour of its flowers.
Some are of a clear white; whilst others are deep
scarlet tinged with a pale pink.

The Indians some apply the decoction of root
to their eyes; and grow here in that state. This species
succeeds best in a shady situation; and in the sub-
mer, should be supplied with copious watering.

From Kew, 2, 1808

CAMPANULA PU'MILA.

DWARF BELL-FLOWER.

Class.
PENTANDRIA.

Order.
MONOGYNIA.

Natural Order.
CAMPANULACEÆ.

Native of Switzerland	Height. 3 inches.	Flowers in June, Aug.	Duration. Perennial.	Introduction not known.
--------------------------	----------------------	--------------------------	-------------------------	----------------------------

No. 48.

Campanula, from the Latin; the diminutive of campana, a bell; a name appropriately bestowed on this genus, from the shape of the flowers. Pumila, also from the Latin, signifying little.

This very pretty campanula has been noticed by the greater part of our botanical writers as a variety of Campanula rotundifolia. It is, however, very different in several particulars, and may well claim the distinction of a separate species.

There are two varieties, the blue and the white; and where the soil is rich and loose, it is difficult to keep their thread-like roots within proper limits; which, if not attended to, may occasion inconvenience, by their mingling with other plants.

During a part of the summer, this plant affords a very attractive border. Its little pendent blue, or more exquisitely delicate white, blossoms, which are yielded in "lovely profusion," form an unbroken line of neatness and simplicity. This effect is best obtained by enclosing the roots between two rows of tiles or slates, placed in the ground edgeways, about two inches apart, with their upper edges even with the surface of the soil.

Hort. Kew. 2, v. 1, 345.

CAMPANULA PUMILA.

SWISS BELL-FLOWER.

Order.
NIGHTSHADE.

Class.
TETRADELM.

Native of Europe.
CAMPANULACEAE.

Native of	Height	Flowers in	Location	Intensification
Switzerland	2 feet	June-Aug.	Swiss Alps	not known

No. 42.

Campanula, from the Latin; the diminutive of *campana*, a bell; a name apparently bestowed on this genus from the shape of the flower. *Pumila*, also from the Latin, signifying little.

This very pretty *campanula* has been noticed by the greater part of our botanical writers as a variety of *Campanula rotundifolia*. It is, however, very different in several particulars, and may well claim the distinction of a separate species.

There are two varieties, the blue and the white; and where the soil is rich and loose, it is difficult to keep their thread-like roots within proper limits; which, if not attended to, may occasion inconvenience, by their mingling with other plants.

During a part of the summer, this plant affords a very attractive border. Its little pendulous blue, or more exquisitely delicate white, flowers, which are yielded in "lovely profusion," form an undulating line of sweetness and simplicity. This effect is best obtained by enclosing the roots between two rows of like or alike, placed in the ground obliquely, about two inches apart, with their upper edges even with the surface of the soil.

Flora Kew. 2, v. 1, 245.





Pyrus Japonica.

13



Genista sagittalis.

14



Gentiana acaulis.

15



Ledum buxifolium.

PYRUS JAPONICA.

JAPAN APPLE-TREE.

Class.
ICOSANDRIA.

Order.
PENTAGYNIA.

Natural Order.
ROSACEÆ.

Native of Japan.	Height. 6 feet.	Flowers in Jan. June.	Duration. Perennial.	Introduced in 1796.
---------------------	--------------------	--------------------------	-------------------------	------------------------

No. 49.

With the exposition and inferences of Linneus before us respecting the word *Pyrus*, as given by Sir J. E. Smith, we cannot admit the deduction of the term from the Greek *PUR*, fire, which is stated to have been given in consequence of the fruit drawing to a point like a flame. De Theis says, from the Celtic *PEREN* comes the Anglo-Saxon *PERE*, the English *PEAR*, and the French *POIRE*. Hence *Pyrus* may easily be formed. According to the same writer, *API*, the Celtic name of a fruit of the same kind, is the origin of the Greek *APIOS*, the German *APFEL*, and our *APPLE*. Some authors, who will have *Pyrus* to be of Greek extraction, deduce it from *APIOS*, with the addition of an *R*.

That our readers may be prepared to meet new names, they should be informed that *Chænomeles Japonica*, and also *Malus Japonica*, are amongst those lately bestowed on this shrub.

The mere mention of *Pyrus* opens to our view such a field of speculations as it were impossible to pass without notice. We have the *Pyrus malus*, under which name is arranged all the varieties of that valuable fruit the Apple; and *Pyrus communis*

botanically including the numberless sorts of pears, so common amongst us.

We can but regret that many favourite old sorts of these fruits are unavoidably falling to decay. Trees, like animals, grow old and diseased; and it is observed, that every bud or graft of such old tree, when attached to another stock, still inherits the age of its original, and its consequent disease. The invigoration yielded by its new alliance is only temporary. Thus, concisely, we warn our readers from vain attempts to propagate healthy trees from worn-out varieties. Every man should propagate from seeds, or graft from sorts that are known to have been recently so raised; and fruitful plantations would be the consequence.

The *Pyrus Japonica* is a great acquisition to our gardens, from the beauty, and from the long succession of its flowers. Indeed it is far more easy to name the season of its flowering, than to say when none are produced. Its fruit, which never appears to ripen here, is extremely stony and ungrateful; nevertheless, after having been laid by for a time, it emits a rich fruit-like odoriferous flavour.

It is well calculated for training against trellis work or a wall, and also makes a pretty standard shrub, when spreading from a single stem about two feet in height. It requires no peculiar care, either as regards soil or situation.

The best method of propagation is by layers, which should be put down in the autumn, and they will strike root freely in the following year. Cuttings are sometimes taken. A part of them will grow, but they produce weak plants.

Hort. Kew. 2, v. 3, 209.

GENISTA SAGITTA'LIS.

JOINTED GENISTA.

Class.
DIADELPHIA.

Order.
DECANDRIA.

Natural Order.
LEGUMINOSÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
Germany.	1 foot.	May, June.	Perennial.	in 1758.

No. 50.

It is pretty generally agreed that Genista has been derived from the Latin genu, the knee; but on what account, authors appear less unanimous. Some say in allusion to the bending, or singular connexion of one part of the stem to another; whilst others conceive that it arose from their similar flexibility; or its utility in relieving pain in that joint. Sagittalis, also from the Latin, signifying, of an arrow, perhaps from the leaf having the appearance of the feathered end of an arrow.

The singular formation of this plant will always claim for it a place in the garden of the curious. The leaves are produced one from the end of another; alternately from the upper and under sides, connected by the midrib, which has a partial termination at the end of each.

It may be divided at the root, though a better mode of propagation is from seeds. These should be sown in the autumn, and the plants kept free of weeds during the next summer. In September they may be transplanted to the situation where they are intended to remain. It succeeds in almost any sort soil, or in any aspect.

Hort. Kew. 2, v. 4, 259.

GENTIANA ACAULIS.

GENTIANELLA.

Class.

PENTANDRIA.

Order.

DIGYNIA.

Natural Order.

GENTIANEÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
Alps of Eur.	3 inches.	April, May,	Perennial.	in 1629.

No. 51.

This genus of plants has received its name in honor of Gentius, a king of Illyria, who is said to have discovered one of the species of it. He is also supposed to have experienced its virtues in his army, as a cure for the plague. Acaulis, from the Greek, signifying stemless, which it is in its native alpine situations. Gentianella is formed as the diminutive of Gentian, to characterize a small species.

It cannot but be sincerely regretted by the zealous votaries of Flora, who happen to possess their parterres of beauty within the influence of the smoke of towns, that this interesting plant is so fastidious as to refuse yielding its exquisitely brilliant blue flowers amongst them; though in high situations whether planted in strong or light soil, it generally flowers in great splendour. Pure air has always been deemed indispensable, yet peculiar management may possibly surmount the difficulty; for in situations where it never, or very rarely, produces flowers, we have generally observed it appear healthy, and increase luxuriantly. This would suggest the propriety of planting it in poor soil, for sometimes by checking luxuriance, we obtain, from some other

plants, both flowers and fruit. Upon this principle it is, that various arts are successfully practised on fruit trees; such as curtailing their roots, cutting notches in the larger limbs, and also that of ringing them. The latter method may, with the greatest confidence, be recommended for practice on young free-growing apple or pear trees, which frequently increase their wood too fast to admit of fruit being produced. It may be useful to some of our readers, and shall be briefly stated.

If blossoms have not been usually produced, ringing should be performed after the fall of the leaf. This is called *procuration ringing*. When trees blossom, but fail to bear fruit, it may be done whilst they are in flower. This is called *maturation ringing*; and will induce the production of much finer fruit than would ever be yielded without it. The operation merely consists in taking a ring of the bark entirely off the whole circumference of a large or small branch or limb of the tree. The breadth of the ring, on a luxuriant apple or pear tree, may be a quarter of an inch; but on those of slow growth it should be rather less.

This practice is not so strictly applicable to stone fruits, from the propensity of the trees to gum and canker about the wound. Yet we have successfully practised it on these; taking care to ring such parts as have free growth, making the ring but narrow, and doing it at a period when the sap is most inactive. Thus may fruit trees be rendered productive, and flowering plants ornamental, where neither was the case before art stepped in and forced the efforts of nature.

LEDUM BUXIFOLIUM.

BOX-LEAVED LEDUM.

Class.
DECANDRIA.

Order.
MONOGYNIA.

Natural Order.
RHODORACEÆ.

Native of	Height.	Flowers in	Duration.	Introduced
Carolina.	1 foot.	April, May.	Perennial.	in 1736.

No. 52.

The derivation of the name of this genus, from the Greek LEDON, has been noticed under No. 47. Buxifolium, from buxus, box; and folium, a leaf; in consequence of the resemblance of the foliage of this plant to that of the common box.

A retrospective glance would seem to indicate the metamorphosis of our "Botanic Garden" into an orchard. We, however, never intend being fettered by the limits of a particular subject. The whole vegetable kingdom is before us, and to the extent of our humble capacity we shall lay hold on any part of it that may seem to develop a source of knowledge, either useful to the hands of the practical, or gratifying to the minds of the speculative.

The *Ledum buxifolium* is a beautiful close-growing little evergreen shrub, rarely reaching the height of twelve inches, which should be planted in sandy peat, in a cool and rather shaded situation; and, as far as is possible, where it may have the advantage of pure air. Its chief demand on our care arises in the heat of summer when it should be very frequently watered, or it will be liable to be lost. It is propagated slowly by layers.

Hort. Kew. 2, v. 3, 48.

The description of the shape of this genus, from the Greek *gaster*, has been noticed under No. 47. *Gasterium*, from *gaster*, belly; and *gaster*, a belly; in consequence of the resemblance of the shape of this plant to that of the common box.

A retrospective glance would seem to indicate the metamorphosis of our "Botanic Garden" into an orchard. We, however, never intend being fettered by the limits of a particular subject. The whole vegetable kingdom is before us, and to the extent of our human capacity we shall lay hold on any part of it that may seem to develop a source of knowledge, either useful to the hands of the practical, or extending to the minds of the speculative.

The *Gasterium* *hirsutum* is a beautiful close-growing little evergreen shrub, rarely reaching the height of twelve inches, which should be planted in sandy soil, in a cool and rather shaded situation; and, as far as is possible, where it may have the advantage of partial shade. Its chief demand on our care arises in the best of autumn, when it should be very frequently watered, so that it will be able to do so. It is propagated slowly by layers.





Lillium tigrinum.

1/4



Erica Australis.



Epilobium Dodonæi.



Tagetes patula.

1/2

LILIUM TIGRINUM.

TIGER-SPOTTED LILY.

Class.

HEXANDRIA.

Order.

MONOGYNIA.

Natural Order.

LILIACEÆ.

Native of	Height.	Flowers in	Duration.	Introduced
China.	4 feet.	July, Aug.	Perennial.	in 1804.

No. 53.

Lily, from the Greek LEIRION. The derivation and application of its trivial name are sufficiently evident.

We know of very few plants that excited more general interest than did the Tiger Lily on its introduction to this country. Every one heard of, admired, and resolved on possessing, this Chinese beauty; and in a very short time, from its facility of propagation, the cottager and nobleman boasted alike of its splendour in their borders. Happily, our nature will not admit the continued exertion of these strong feelings of delight, which are generated by novelty; or, we should be unceasingly carried about by ecstasies, and temperate reason could no where build her throne.

In a former number the resemblance of the bulb and the bud was hinted at. Their utility, as regards vegetation, is precisely the same; for they both constitute what Linneus calls the hybernaculum, or the winter quarters, of the young plant. Their principal difference exists in the situation which they occupy; and in the present, and a few other plants, even this distinction is wanting.

‘I consider (says T. A. Knight, Esq. in a letter to the author,) a bulb to differ from an ordinary bud, only in having a reservoir of a different form attached to it. The bud of a tree, or of the tuberous root of a potatoe, is attached to a mass of alburnum and bark, from which, when it germinates, it draws its requisite nutriment. A bulbous root, such as the common onion, has numerous thick and fleshy scales, which in their incipient state, might have extended into leaves, but which, instead of extending themselves, remain short and are distended by becoming reservoirs of the true sap of the plant, as the bark and alburnum were in the cases before-mentioned.’

He observes, also, ‘Buds of every kind have their attached reservoirs, without which they cannot live and extend themselves. Some species of trees and herbaceous plants possess a power which others do not, of re-producing buds upon the surface of their alburnum. It is, however, the unanimous opinion of the continental naturalists, and of the English and Scotch, with the exception of myself, that all buds originate from the Medulla; and it is true that in all cases, almost, a bud may be traced to the Medulla; but I have, in a great many instances, occasioned buds to be generated upon the smooth surface of the alburnum; and I have often seen them thus produced naturally.’

If the bulbs which are produced on the stems of this magnificent Lily, be planted in the borders as soon as they fall, or can be easily detached, they will grow, and in most cases blossom at three years old, but stronger in the fourth year.

Hort. Kew. 2, v. 2, 141.

ERICA AUSTRALIS.

SPANISH HEATH.

Class.
OCTANDRIA.

Order.
MONOGYNIA.

Natural Order.
ERICÆ.

Native of Spain.	Height. 2 feet.	Flowers in April, May.	Duration. Perennial.	Cultivated in 1769.
---------------------	--------------------	---------------------------	-------------------------	------------------------

No. 54.

Erica, from the Greek EREICO; see No. 22. Australis, from the Latin, southern; a specific name given in consequence of its having been introduced from the southern part of Europe.

This plant will always prove a peculiarly interesting appendage to the peat border, and should never be dispensed with. The hardy heaths form a little tribe of shrubs whose beauties we cannot class with the splendour of the Kalmias, the Azaleas, and the Rhododendrons; but they equally interest us through a far different medium. They introduce themselves to our feelings by their modesty and humility; and we readily admit the propriety of Dr. Watts's assertion—

“Humility is a plant of lovely growth.”

Still the humble growth of some of the tender species of Erica, whose flowers are occasionally very specious, may further remind us of him who is humble only to embellish his grandeur.

The Erica australis should be planted in sandy peat; or in a mixture of peat and fresh loam; and like most other of the hardy heaths, though they make root but slowly, may be increased by layers.

Hort. Kew. 2, v, 2, 396.

EPILOBIUM DODONÆI.

DODONÆUS'S EPILOBIUM.

Class.
OCTANDRIA.

Order.
MONOGYNIA.

Natural Order.
ONAGRARIÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
Switzerland.	9 inches.	July, Aug.	Perennial.	in 1798.

No. 55.

Epilobium is compounded of three Greek words, EPI LOBOU ION, a violet upon a pod; or, more literally, upon - a pod - a violet: not that a violet resembles the blossom, but is intended to indicate a beautiful flower. Dodonæi from Dodonæus, an eminent physician and botanist of Friesland, who lived in the sixteenth century, and published several botanical works in German, illustrated by wood cuts, similar to those of Gerard and Parkinson.

This species has been noticed, by some writers, as synonymous with Epilobium angustissimum. Ours is a plant with procumbent stems, and otherwise differing from angustissimum.

It is the prettiest plant that we know of the tribe, and is never troublesome by spreading at the root, as are some species of Epilobium.

After it has done blossoming, the whole of the stems may be cut off, or they will continue to grow, and thus, sometimes, lessen the vigour of the roots. It is easily increased by separating the young shoots in the spring, which will succeed, notwithstanding they may be entirely devoid of any fibrous appendage.

Richenbach Icones, 4, 342.

EPILOBION DROSERAE

ROSEMARY'S EPILOBION

Class.
Rosaceae

Class.
Rosaceae

Number of
leaves - 21

Number of leaves	Height of plant	Flowers in leaf axils	Flowers in leaf axils
21	1.5 m	1.5 m	1.5 m

No. 21

Epilobion is composed of three Greek words, *epi* upon, *ros* a rose, and *bion* a life; it is a plant which lives in the sixteenth century, and published several botanical works in German, illustrated by wood cuts, similar to those of Gerard and Parkinson.

This species has been noticed by some writers, as synonymous with *Epilobium angustifolium*. Ours is a plant with succulent stems, and other, also differing from *angustifolium*.

It is the earliest plant that we know in the north, and is never troublesome by spreading at the root, as are some species of *Epilobium*.

After it has done its duty, the whole of the stems may be cut off, or they will continue to grow, and then, sometimes, form the vigour of the roots. It is easily increased by separating the young shoots in the spring, which will succeed, notwithstanding they may be entirely devoid of any fibrous appendages.

Richardson, June 4, 1858

TAGE/TES PAT'ULA.

FRENCH MARYGOLD.

Class.
SYNGENESIA.

Order.
POLYGAMIA SUPERFLUA.

Natural Order.
CORYMBIFERÆ.

Native of Mexico.	Height. 3 feet.	Flowers in Aug. Oct.	Duration. Annual.	Cultivated in 1596.
----------------------	--------------------	-------------------------	----------------------	------------------------


No. 56.

Apuleius, a celebrated platonic philosopher of the second century, is said to have used this term; its derivation must therefore be doubtful. De Theis has derived it from Tages, an Etruscan deity, grandson of Jupiter, and teacher of divination. Patula, from the Latin, spreading.

Why this plant is called French Marygold is now somewhat difficult to determine; but it is more than probable that it received this appellation in consequence of its seed having been first imported to this country from France.

Its cultivation is so generally known, that nothing need be said respecting it; except to warn our readers against a formidable enemy to the young plants. If they be much eaten, a single examination, late at night, with the assistance of a light, will show the depredators to be young earwigs, (*Forficula auricularia*). Woollen cloth, loosely folded; hollow bean stalks; or two small boards, placed upon each other, with one edge of the upper one raised sufficiently to admit their creeping between them, will form useful traps, and the insects may be destroyed every morning.

Hort. Kew. 2, v. 5, 88.





Dracocephalum speciosum.

$\frac{1}{2}$



Buddlea globosa.

$\frac{1}{2}$



Phlox ovata.

$\frac{2}{3}$



Primula acaulis.

$\frac{2}{3}$

DRACOCEPH'ALUM SPECIO'SUM.

SHOWY DRAGON'S HEAD.

Class.
DIDYNAMIA.

Order.
GYMNOSPERMIA.

Natural Order.
LABIATÆ.

Native of N. America.	Height. 3 feet.	Flowers in August.	Duration. Perennial.	Introduced in 1822.
--------------------------	--------------------	-----------------------	-------------------------	------------------------

No. 57.

Dracocephalum is a term compounded of two Greek words, DRACON, a dragon; and CEPHALE, the head, on account of the fancied resemblance of its corolla to the head of that fabulous animal. Speciosum, from the Latin, showy.

This is a handsome erect herbaceous plant, of somewhat larger growth than the Dracocephalum denticulatum, to which it bears considerable resemblance, but from its late introduction, is not so frequently met with.

It will grow in any common garden soil, but prefers a rich and rather light loam. It increases freely at the root, and may be divided in spring, or in the autumn. If required, cuttings of the young stems may be struck for increase.

We ought not, perhaps, to pass by the present opportunity of making further mention of the ideal animal whose name has assisted in distinguishing this genus of plants. We say ideal, because much that we have heard and read of dragons, during childhood, must have been so. Such tales may serve the purpose of the nurse, but are highly ridiculous, and improper to be implanted in the youthful mind.

No opportunity should, therefore, be lost in undeceiving the juvenile intellect, and informing it that those tales of flying monsters vomiting fire, the glance of whose eyes is death to the beholder, have wholly originated in the glowing imaginations of ancient mythologists and poets. It is true that we have frequent mention of dragons in the sacred writings; but there the term, as Bishop Lowth observes, is used allegorically for great potentates,—enemies and persecutors of the people of God.

Linneus has, in some degree, aided in perpetuating those tales of wonder, by applying the name draco, or dragon, to distinguish a genus of amphibious animals of the reptile kind; which he conceived bore some resemblance to the fancied form of the dragon of the ancients. One species of it is aptly enough called draco volans, or flying dragon, being furnished with wing-like membranes, which enable it to follow its insect prey a short distance through the air. It is an inhabitant of Africa and Asia; but instead of the very formidable monster which the imagination may be expected to create, it is merely an inoffensive Lizard, of about ten inches in length.

Several accounts of dragons have been circulated amongst us, through mediums of respectability. One of these is given in the *Encyclopædia Britannica*, with a plate of the animal itself. An account of the same creature appeared in the *Gentleman's Magazine* for 1749; but this was discovered to be a ridiculous cheat; the specimen shown being more nor less than the skin of a species of shark, which had been distorted and dried, to impose on the credulous.

Sweet's Fl. Gard. 93.

BUD'DLEA GLOBO'SA.

ROUND-HEADED BUDDLEA.

Class.
TETRANDRIA.

Order.
MONOGYNIA.

Natural Order.
SCROPHULARINÆ.

Native of	Height.	Flowers in	Duration.	Introduced
Chili.	10 feet.	May, June.	Perennial.	in 1774.

No. 58.

The appellation Buddlea was bestowed on this genus of plants by Dr. Houston, in honour of Adam Buddle, an English botanist, of some note in the 17th century. He formed an herbarium, which is now deposited in the British Museum. Globosa, from the Latin, round.

The flowers of this species of Buddlea are interesting, inasmuch as they differ materially from those of the usual garden shrubs. They possess a peculiar fragrance, approaching to that of honeycomb; on which account, and from the compact mode of its inflorescence, it is sometimes called the honeycomb plant.

Its habit of growth, and the southern latitude of its native climate, do not qualify it for a standard in our shrubberies; but when trained against a wall, of southern aspect, it will grow luxuriantly and produce abundance of flowers.

It may be propagated by cuttings, taken about midsummer, which should be planted under a hand-glass, on a hotbed, if convenient. Young plants must be protected from frost; and a slight covering, in severe weather, will benefit old ones.

Hort. Kew. 2. v. 1, 250.

PHLOX OVATA.

OVAL-LEAVED PHLOX.

Class.
PENTANDRIA.

Order.
MONOGYNIA.

Natural Order.
POLEMONIACEÆ.

Native of N. America.	Height. 6 inches.	Flowers in May, June.	Duration. Perennial.	Cultivated in 1790.
--------------------------	----------------------	--------------------------	-------------------------	------------------------

No. 59.

Though the term *Lychnidea* has been expelled from Botanical genera, it is sometimes, nevertheless, used as an English name for this tribe of plants; see No. 6. *Ovata*, from the Latin, signifying egg-shaped, or of an oval figure.

The *Phlox ovata* is a beautiful herbaceous plant, and however often met it will always be welcome. We believe there is not one individual in this tribe but has powerful claims to the attention of every true florist; we may therefore presume that our readers will have pleasure in being occasionally introduced to others of the same family.

This little plant, though from North America, was for many years after its introduction, regarded as a subject for the greenhouse, and indeed in moist situations it will occasionally decay.

The greater part of the *Phloxes* delight in peat, or a mixture of that soil with loam; peat, however, seems rather too retentive of moisture to suit the *Phlox ovata*; but if planted in sandy loam, in a warm situation, the severest frosts will not be found to injure it. It may be propagated by parting the roots, or by cuttings.

Hort. Kew. 2, v 1. 327.

PRIMULA ACAULIS;

Flore pleno rubra, et flore pleno alba.

PRIMROSE;

Double Crimson, and Double White.

Class.

PENTANDRIA.

Order.

MONOGYNIA.

Natural Order.

PRIMULACEÆ.

Native of Britain.	Height. 3 inches.	Flowers in April, May.	Duration. Perennial.	Cultivated in 1640.
-----------------------	----------------------	---------------------------	-------------------------	------------------------

No. 60.

This plant has received its name from the Latin *primus*, first, on account of its early flowering; though not quite correctly, for we have aconites, snowdrops, and crocuses, before the delicate primrose dares to unfold its beautiful petals. *Acaulis*, from the Greek, stemless.

The beautiful varieties of primroses which inhabit our gardens, whether crimson, lilac, or white, double or single, are alike included under one species. Indeed, much difficulty has arisen respecting the origin, and consequently, the proper specific division, of several of the *primula* tribe. Linneus considers the primrose, the cowslip, and oxlip, originally but one distinct species, and several reasons may be advanced in support of this theory; yet still the partially distinct characters of the primrose and cowslip should incline us now to separate them, whatever may have been their origin. We certainly have seen flowers of the primrose supported on a scape or stalk, and thus approach the oxlip; whilst the cowslip, when brought into cultivation, will have its flowers enlarged, and thus also incline towards the oxlip. Hence a tendency is shown, in the two ex-

tremes of distinction, to verge towards each other; indeed, a host of connecting links between these plants will present themselves to the diligent and inquiring botanist.

Be the scientific difference or connection of these British subjects whatever it may, it does not lessen the value of our attractive and modest primrose, that lives on banks and hides its beauties beneath the brambles' shade,—

“Lorn tenant of the peaceful glade,
Emblem of virtue in the shade,
Rearing thy head to brave the storm
That would thy innocence deform.
Of all the flowers that greet the spring,
Of all the flowers the seasons bring;
To me while doom'd to linger here,
The lowly Primrose shall be dear.” MAYNE.

The two peculiarly pleasing varieties of which we now present figures are the most elegant little subjects that we are acquainted with in the species. They are usually planted in a loamy soil; we, however, find a sandy peat, with a little loam, more suitable to their growth. As the double white does not freely produce offsets that can be conveniently slipped from the old plant, it will be found an advantageous practice to slit the thick part of the old root longitudinally with a knife, into as many parts as the head will admit; observing to retain a portion of the fibrous roots to each division. These being planted in pots or the borders, should be regularly supplied with water till they have taken root; but during the winter, if in pots, they require little or no water.


Hort. Kew. 2, v. 1, 307.

LOVELLIA, 1878

PLANT. SPERMATOPHYTES.



The plant is a large, upright, and
erect, with a single, or a few, stems
arising from a common base, and
bearing a large, flat-topped, or
cymose, cluster of flowers at the
summit. The leaves are compound,
and are borne on long, slender
petioles. The flowers are large,
and are borne in a loose, cymose,
cluster at the summit of the stem.





Lobelia syphilitica.

$\frac{1}{2}$



Coreopsis tenuifolia.

$\frac{1}{2}$



Rhododendron hirsutum



Cochlearia Greenlandica.

LOBELIA SIPHILITICA.

BLUE LOBELIA.

Class.
PENTANDRIA.

Order.
MONOGYNIA.

Natural Order.
CAMPANULACEÆ.

Native of Virginia.	Height. 18 inches.	Flowers in Aug. Sept.	Duration. Perennial.	Cultivated in 1665.
------------------------	-----------------------	--------------------------	-------------------------	------------------------

No. 61.

Lobelia is a name instituted by Plumier, after Matthias de Lobel, a Flemish botanist of the sixteenth century; who, in youth, acquired an ardent love of plants; and, through life, cultivated the science of botany with considerable success. He was appointed botanist to King James I., and died near London, at the advanced age of seventy-eight. Woodville observes, that this plant derived its appellation, Siphilitica, from its efficacy in the cure of siphilis, according to the experience of the North American Indians. As its antisiphilitic powers have not, however, been confirmed by European practice, it may be needless to treat at all on its medical qualities.

Several plants of the Lobelia tribe possess very active medicinal properties, particularly the Lobelia Tupa, a native of Chili. This species is poisonous in the extreme, and acts as an emetic, simply by smelling the flowers.

Whether the Lobelia siphilitica be planted in the open ground, or kept in a pot, the offsets should be divided in the spring; and they will blossom freely without further care.

LOBELIA SIPHILITICA.

RED LOBELIA.

General Description.

Height.	Flowers in	Duration of
1 to 2 feet.	terminal.	in fruit.

No. 41.

Lobelia is a native introduced by the Spaniards. Matthias de Lobel, a Flemish botanist of the sixteenth century; who, in 1594, accompanied an expedition to Florida; and, through his exertions, the first seeds of this plant were introduced into Europe. He was appointed physician to King James I., and died near London, at the advanced age of seventy-eight. *Wendland* observes, that the plant derived its specific name, *siphilitica*, from its use in the cure of syphilis, according to the experience of the North American Indians. As its antisyphilitic powers have not, however, been confirmed by European practice, it may be needless to treat at all on its medicinal qualities.

Several kinds of the Lobelia *siphilitica* are active medicinal properties, particularly the *Lobelia inflata*, a native of North America. This species is common in the United States, and acts as an emetic, chiefly by stimulating the bowels.

When the Lobelia *siphilitica* is planted in the open ground, or kept in a pot, the plants should be divided in the spring; and they will be ready for use in the autumn.

How to use it. 1. 1830.

COREOPSIS TENUIFOLIA.

SLENDER-LEAVED COREOPSIS.

Class. SYNGENESIA. *Order.* POLYGAMIA FRUSTRANEA.
Natural Order. CORYMBIFERÆ.

Native of N. America.	Height. 15 inches.	Flowers in July, Aug.	Duration. Perennial.	Introduced in 1784.
--------------------------	-----------------------	--------------------------	-------------------------	------------------------

No. 62.

Coreopsis, from the Greek CORIS, a bug; and OPSIS, appearance; in allusion to the apparent similarity of the seed and the insect. From this circumstance it is, in part, that these plants are sometimes called tick-seed sunflower. Tenuifolia, from the Latin tenuis, slender; and folium, a leaf.

There were formerly between thirty and forty distinct species of Coreopsis described; and nearly the whole of them, like tenuifolia, hardy, and herbaceous. The genus has, however, been somewhat curtailed, from the generic characters of some of its former species having been found such, as of necessity, required their removal to other genera. They are generally showy plants; and more perfect instances of their attraction need not be given, than in the two species already published in this work, No. 13, and No. 46. The former of these, the Coreopsis tinctoria, has spread over Great Britain more rapidly than any plant we have ever known. In the three years, since the period of its introduction, its beauty has secured it a passport to almost every respectable garden in the kingdom; where it will continue to be cultivated as one of the most

desirable annuals that our transatlantic friends have ever bestowed upon us.

The *Coreopsis tenuifolia* requires no peculiar care, but may be planted in any common garden soil. Its increase at the root will admit a division every other year, or even more frequently.

It is a desirable plant, from the compactness of its growth and neatness of its slender foliage. It opens its brilliant flowers before the gay annuals of the autumn are over prominent; a season which is usually burthened with these tints of gold. Nature, indeed, seems prodigal of this rich dress. The opening of spring exhibits her in the garden, conspicuous in aconites, crocuses, and the gay variety of narcissuses; whilst the meadows, in a blaze of buttercups and cowslips, remind us of the descent of Jupiter in a shower of gold.

The pleasures of hope are multiplied as we approach the object of anticipation; and now, at the vernal equinox, the very mention of spring is exhilarating in the extreme, when

“ All that is sweet to smell, all that can charm
Or eye or ear, burst forth on every side,
And crowds upon the senses.

“ By nature's swift and secret working hand,
The garden glows, and fills the liberal air
With lavish fragrance; while the promised fruit
Lies yet a little embryo, unperceived,
Within its crimson folds.

“ Ye fostering breezes, blow,
Ye softening dews, ye tender showers, descend!
And temper all, thou world-reviving sun,
Into the perfect year.”

THOMSON.

Hort. Kew. 2, v. 5, 133.

RHODODENDRON HIRSUTUM.

HAIRY-LEAVED RHODODENDRON.

Class.
DECANDRIA.

Order.
MONOGYNIA.

Natural Order.
RHODORACEÆ.

Native of Switzerland.	Height. 18 inches.	Flowers in May, July.	Duration. Perennial.	Introduced in 1739.
---------------------------	-----------------------	--------------------------	-------------------------	------------------------

No. 63.

Rhododendron is deduced from two Greek words, **RODON**, a rose, and **DENDRON**, a tree. The name was first adopted by Dioscorides. Linneus's application of it does not, however, appear to be continued to the same plants. Our Nerium is supposed to have first borne the appellation. Hirsutum, from the Latin, rough or hairy.

Our present subject bears a strong affinity to the Rhododendron ferrugineum, excepting that the under side of its leaves are less rusty, and they are also fringed with rigid hairs, which are not found on the leaves of the Rhododendron ferrugineum.

It is rather more free in growth than the last-mentioned species, and generally possesses an advantage over it, in the abundance of its beautiful blossoms.

These plants are propagated, in Great Britain, principally by layers; as they rarely produce seeds or suckers, except in their natural climate. The usual time of laying them, is the latter part of summer, and the heads only of the young shoots should be left above the soil. Peat soil and a northerly aspect will be found most suitable.

Hort. Kew. 2, v. 3, 49.

COCHLEARIA GÆNLANDICA.

GREENLAND SCURVY-GRASS.

Class.
TETRADYNAMIA.

Order.
SILICULOSÆ.

Natural Order.
CRUCIFERÆ.

Native of Greenland.	Height. 2 inches.	Flowers in June.	Duration. Perennial.	Inhabits High Hills.
-------------------------	----------------------	---------------------	-------------------------	-------------------------

No. 64.

From the Latin cochleare, a spoon ; a term applied to this family of plants from the formation of their leaves being concave, and resembling an old fashioned spoon. Grœnlandica, from the country where it has been found. It has, occasionally, been met with also in the Orkneys, and on the mountains of Scotland.

This species is, by some authors, termed a starved variety of the officinalis, an English species, pretty well known in the north, and on the sea coast, a plant which has obtained for the genus the title of scurvy-grass. Its efficacy in scorbutic affections appears to be established on the most respectable authorities, and though various preparations of the *Cochlearia officinalis* are prescribed, it is generally acknowledged that the green plant taken as a salad, is by far the most efficacious mode of employing it as an antiscorbutic.

The best method of keeping the Greenland scurvy-grass, is in a small pot of light loam ; and like most other alpine plants, it succeeds best in a high and open situation, where it has the full advantage of pure air.

Withering, 678.

COCHLEARIA GEMMATA.

GREENLAND SCURVY-GRASS.

Order
GROSSULARIACEAE

Family
GROSSULARIACEAE

Native of
Greenland

Native of	Height	Flowers in	Parentage	Localities
Greenland	2 inches	June	Parental	High Hills

No. 64

From the Latin cochlear, a spoon; a term applied to this family of plants from the formation of their leaves being concave, and resembling an old fashioned spoon. Gravelled, from the country where it has been found. It has, occasionally, been met with also in the Orkneys, and on the mountains of Scotland.

This species is, by some authors, termed a stunted variety of the officinalis, an English species, pretty well known in the north, and on the sea coast, a plant which has obtained for the genus the title of scurvy-grass. Its efficacy in scurvy affections appears to be established on the most respectable authorities, and though various preparations of the Cochlearia officinalis are prescribed, it is generally acknowledged that the green plant taken as a salad, is by far the most efficacious mode of employing it as an antiscorbutic.

The best method of keeping the Greenland scurvy-grass, is in a small pot of light beer; and like most other alpine plants, it succeeds best in a high and open situation, where it has the full advantage of pure air.

W. H. W.

PLANTING GUIDE

1881



These plants are very hardy and will grow in any soil.

They are very useful for many purposes.

They are very hardy and will grow in any soil.

They are very useful for many purposes.

They are very hardy and will grow in any soil.

They are very useful for many purposes.

They are very hardy and will grow in any soil.

They are very useful for many purposes.

They are very hardy and will grow in any soil.

They are very useful for many purposes.

They are very hardy and will grow in any soil.

They are very useful for many purposes.

They are very hardy and will grow in any soil.


They are very useful for many purposes.

They are very hardy and will grow in any soil.

They are very useful for many purposes.

They are very hardy and will grow in any soil.

They are very useful for many purposes.



These plants are very hardy and will grow in any soil.

They are very useful for many purposes.



Dianthus Chinensis.

½



Sisyrinchium striatum.

½



Calceolaria corymbosa.



Philox. cetacea.

½

DIANTHUS CHINENSIS.

CHINA PINK.

Class.
DECANDRIA.

Order.
DIGYNIA.

Natural Order.
CARYOPHYLLÆ.

Native of China.	Height. 9 inches.	Flowers in July, Sep.	Duration. Perennial.	Cultivated in 1713.
---------------------	----------------------	--------------------------	-------------------------	------------------------

No. 65.

Linneus, duly appreciating the beauty and fragrance of this genus of plants, seems to have been desirous of peculiarly distinguishing it by a name, and called it Jove's Flower; deriving Dianthus from the Greek DIOS, of Jove, and ANTHOS, a flower. Chinensis, from its native country.

This plant is usually mentioned as biennial. We have considered it perennial, from conviction that it most probably is perfectly so in its native climate. It may be cultivated with advantage as an annual, as it perfects its seed in the first season of its growth, but having found it continue three years in a healthy state, it would be improper to term it annual or biennial.

The numerous combinations of colour into which this well-known little subject is prone to sport, renders it truly interesting. Each succeeding flower may be anticipated as more beautiful than its predecessor.

The China Pink, or Indian Pink, as it is sometimes called, may be raised from seeds, which should be sown on the borders, or a hot-bed, at the latter end of March. A dry light soil is best suited to it.

DIANTHUS CHIRSIOPSIS

CHINA

Color

Flowers

Stems and leaves

Characteristics

Height	Flowers in	Flowers in
1 foot	10 to 15	10 to 15

No. 10.

This plant is usually mentioned as biennial. It has been considered as perennial from observation that it most probably is perennating in its native climate. It may be cultivated with advantage as an annual, as it produces its seed in the first season of its growth, but having found it continues three years in a healthy state, it would be improper to state it as annual or biennial.

The numerous combinations of colors, from which this well-known little subject is known to sport, are here in high perfection. Each successive flower may be cultivated as a new beautiful form in the garden.

This plant is called in Indian 'Lok', and in Chinese 'Lok'. It is also called 'Lok' in some parts of China, and 'Lok' in some parts of Japan.

This plant is called in Indian 'Lok', and in Chinese 'Lok'. It is also called 'Lok' in some parts of China, and 'Lok' in some parts of Japan.

From the A. J. 1847.

SISYRINCHIUM STRIATUM.

STREAKED SISYRINCHIUM.

Class.
TRIANDRIA.

Order.
MONOGYNIA.

Natural Order.
IRIDÆ.

Native of Mexico.	Height. 18 inches.	Flowers in June, July.	Duration. Perennial.	Cultivated in 1788.
----------------------	-----------------------	---------------------------	-------------------------	------------------------

No. 66.

The word *Sisyrinchium* is deduced from the Greek *sus*, a hog, and *rygchos*, a snout. This name has been handed down to us from classic authors, who lived before the birth of Christ; but it has not been precisely determined what plant then bore the appellation. There is nothing in this genus that entitles it to such a name; nothing, at least, that we can trace. *Striatum*, from the Latin, in reference to its striated leaves and flowers.

In the late general catalogues, this plant is called *Marica striata*, after Curtis; but we follow the authority of the most eminent modern authors and nurserymen, by retaining its previous appellation.

The *Sisyrinchium striatum* is well adapted for the mingled flower border of herbaceous plants, as it continues in bloom during the whole of June and July, and exhibits a pretty variety of flowers in connection with its Iris-like foliage.

It is easily increased, by a division of its roots in autumn or spring, and should be planted in a strong loamy soil. As it is sometimes destroyed by severe frosts, it will be advisable to protect a plant in the cold frame.

Hort. Kew. 2, v. 4, 136.

STYRACHEA STYRACIUM

STYRACIUM STYRACIUM

China
Yunnan
Natural Order
Rosaceae

Native of	Height	Flowers in	Domestic	Collected
Mexico	12 inches	June, July	Present	in the

No. 66

The word *Styracium* is derived from the Greek *styrax*, a bog, and *styrax*, a resin. This name has been handed down to us from classic authors, who lived before the birth of Christ; but it has not been precisely determined what plant they bore the appellation. There is nothing in the genus that entitles it to such a name; neither, at least, that we can trace. *Styracium* from the Latin, in reference to the stinked leaves and flowers.

In the late general catalogue, this plant is called *Styracis styracis*, after Eurtius; but we follow the authority of the most eminent modern authors and our own, by retaining its previous appellation.

The *Styracium styracis* is well adapted for the mingled flower border or herbaceous plant, as it continues in bloom during the whole of June and July, and exhibits a pretty variety of flowers in connection with its luscious foliage.

It is easily increased, by a division of its roots in autumn or spring, and should be planted in a strong loamy soil. As it is sometimes destroyed by winter frosts, it will be advisable to protect a plant in the cold frame.

Bot. Kew. 3. v. 156.

CALCEOLARIA CORYMBO'SA.

CHILI SLIPPER-WORT.

Class.
DECANDRIA.

Order.
MONOGYNIA.

Natural Order.
SCROPHULARINÆ.

Native of	Height.	Flowers in	Duration.	Introduced
Chili.	12 inches.	June, July.	Annual.	in 1823.

No. 67.

The term, Calceolaria, has originated from the Latin calceolus, a little shoe : and an inspection of the figure of the plant will sufficiently show the reason of its bearing the appellation. Corymbosa, from its mode of inflorescence.

The shape of the corolla of Calceolaria is one of those which, from its infrequency, attracts the notice of the most careless observer. It cannot be said to present any phenomenon that does not exist in the simplest flower that we meet with. Its novelty alone surprises. Were we accustomed to see none but monopetalous flowers, similar to the present one, how excessive would be our surprise and pleasure on first beholding the brilliantly rayed daisy, with its golden engine-turned centre; which now is pressed beneath our feet, and regarded, almost as little by the botanist of sensibility, as by the rudest hind that ever despoiled its beauties.

The cultivation of the Calceolaria corymbosa is by no means difficult. It may be planted in the open borders, or kept in a pot of loam and peat, but should have a little protection against the severe frosts of our winters.

Bot. Reg. 723.

CAECALIA CORYMBOSA

CHILLI ALPINE-WORT

CHILLI
ALPINE-WORT

CHILLI
ALPINE-WORT

CHILLI
ALPINE-WORT

Native of Chilli	Height 12 inches	Flowers in June, July	Flowers in June, July	Introduced in 1852
---------------------	---------------------	--------------------------	--------------------------	-----------------------

No. 17

The term, *Caecalia*, has originated from the Latin *caecus*, a little obscure; and an inspection of the figure of the plant will sufficiently show the reason of its bearing the appellation. *Corymbosa*, from its mode of inflorescence.

The shape of the corolla of *Caecalia* is one of those which, from its infrequency, attracts the notice of the most careless observer. It cannot be said to present any phenomenon that does not exist in the simplest flower that we meet with. Its novelty alone surprises. When we are accustomed to see those but monophyllous flowers, similar to the present one, how excessive would be our surprise and pleasure on first beholding the brilliantly rayed disk, with its golden saffron-tinted centre; which now is pressed beneath our feet, and regarded, almost as little by the botanist of sensibility, as by the rustiest kind that ever despoiled its beauties.

The cultivation of the *Caecalia corymbosa* is by no means difficult. It may be planted in the open borders, or kept in a pot of loam and peat, but should have a little protection against the severe frosts of our winter.

See No. 17

PHLOX SETACEA.

FINE-LEAVED PHLOX.

Class.
PENTANDRIA.

Order.
MONOGYNIA.

Natural Order.
POLEMONIACEÆ.

Native of	Height.	Flowers in	Duration.	Introduced
N. America.	3 inches.	April, May.	Perennial.	in 1786.

No. 68.

The probable origin of the word Phlox has been noticed under the sixth subject given in this work. Setacea, from the Latin seta, a bristle, in allusion to its bristle-like foliage.


The species of Phlox which we now present to our readers, is completely dissimilar from either of those already treated of, but is still a desirable one. It is a partial trailer, and any effort to lead it upright will rarely be found to improve its effect. Many plants require our care to support them against the assaults of rude winds and battering rains. Our convenience also requires the cutting back, and the confinement of plants in certain positions, otherwise nature, left to herself, generally produces a freedom of outline that must abash every advocate of antique Italian clipping and carving.

We beg to warn those of our fair readers, who happen not to examine the present plant botanically, against confounding it with the Phlox subulata, which is far more common through the midland counties of England. The leaves of the Phlox setacea are narrower than those of subulata, and its flower also possesses greater delicacy of colouring,

not having so dark a centre as the similar species. The *Phlox setacea* has, with us, withstood the severity of several winters in an exposed situation. This circumstance may be adduced as evidence of its hardy nature. It will, however, be occasionally lost, which accounts for its not being more common. A few cuttings, therefore, should be planted in a pot, in June, and covered over with a small bell glass. These should be watered whenever the soil becomes dry, and they will strike root without the assistance of artificial heat. The glass should be taken off for half an hour, three or four times a week, whilst in the shade, and may be entirely removed at the end of six weeks. Plants propagated in this way may, with convenience, be placed for protection, during the winter, within a cold frame. This indulgence will, however, render them somewhat more susceptible of cold than those which have weathered the severity of the season; and if they be turned into the borders before flowering, the keen winds which sometimes occur in March and April will partially dry up the juices of the plant, and render it brown and unsightly.

The slugs are determined enemies of this plant, and great vigilance is necessary to protect it from their depredations. They eat out the tender ends of the shoots, and disappoint our hopes of a luxuriant blossom. Complete protection from these invaders of our pleasures is sometimes rather difficult; if, however, large leaves, or other vegetable refuse, of almost any description, under which they can easily retreat, be laid near to their haunts, they may generally be captured.

Hort. Kew. 2, v. 1, 327.





Fumaria nobilis.

7/8



Hesperis matronalis.

1/2



Symphytum Bohemianum.

7/8



Daphne gnidium.

FUMARIA NOBILIS.

GREAT-FLOWERED FUMATORY.

Class.
DIADELPHIA.

Order.
HEXANDRIA.

Natural Order.
PAPAVERACEÆ.

Native of Siberia.	Height. 6 inches.	Flowers in April, May	Duration. Perennial.	Introduced in 1783.
-----------------------	----------------------	--------------------------	-------------------------	------------------------

No. 69.

The term *Fumaria* is allowed to have had its origin in the Latin *fumus*, smoke; but on what account is not so easy to determine. It has been said, from its affecting the eyes like smoke. This, however, after examining several species, we cannot corroborate. Some of the old herbalists call it smoke of the ground; from which it may, possibly, be inferred, that they named it from its glaucous, or smoke-like appearance, when viewed in considerable quantity, at a distance. *Nobilis*, from the Latin, signifying noble or excellent.

This plant is desirable for its very hardy nature, as well as its gaiety at that particular period of spring, when a blank is wont to pervade our borders: when we have seen the crocuses, hepaticas, scillas, and earlier beauties pass away, and but few of their successors bold enough to venture forth.

It will flourish in any light garden soil, though with but little increase. Its seeds are but sparingly perfected in England, therefore its propagation must principally depend on offsets; which may be separated at any time after the decay of its leaves. Transplanting weakens the roots.

HESPERIS MATRONALIS. *purpurea plena.*

DOUBLE-PURPLE ROCKET.

Class.
TETRADYNAMIA.

Order.
SILIKUOSA.

Natural Order.
CRUCIFERÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
Italy.	30 inches.	June, Aug.	Pereannal.	in 1597.

No. 70.

The name of this genus has been considered under the white variety of the same species, No. 39. We should not, indeed, have noticed the present variety of the same flower, but on account of its very distinct habits from the double white Rocket. It grows much taller, branched, and its flowers are not so closely clustered.

It is suitable for the front of the shrubbery border, and may be placed in those situations where it will be succeeded by annuals of middle growth.

The double varieties of the Rocket, particularly the white, require some attention, in order to the attainment of strong and luxuriant blossoming plants: for we have not only seen them continue to grow in a weakly and unthriving state, but, not unfrequently, die altogether.

The source of the best practical instructions on this head has been previously noticed; and the greatest service we can render our readers will be in transcribing it.

“In the root method of proceeding, with the intention of providing offsets more abundantly, some of the best plants should be placed in an open bed or

border, and not suffered to run up fully to flower; but as soon as the flower stems have advanced eight or ten inches in height, cut them down as close to the ground as possible; and as they shoot again to have them also cut off; for by stopping their upright growth in this manner, the roots are induced more readily to throw out young offsets from their sides, which will be well formed by the beginning of the autumn, when the whole root should be taken up, and the offsets separated from it, and planted out in a nursery bed at about six inches distance, in order to continue until the beginning of autumn, or the following spring; at either of which times they should be carefully removed, with good balls of earth about their roots, to the places where they are to grow for flowering. This mode of treatment will be found peculiarly advantageous.

Such of the flower stems as are thus cut down occasionally, for the purpose of increasing the number of offsets, may be formed into cuttings of proper lengths, and planted out in a shady border, depositing them two parts within the ground, and about three inches assunder, water being given at the time, and repeated as there may be occasion.

In most cases, a number of the cuttings will have stricken good root, and formed shoots at the tops in the course of six or eight weeks. But in order to promote their taking root, in a more effectual manner, they ought to be covered closely with bell or hand-glasses as soon as they are planted, raising them occasionally as the plants begin to shoot at the tops, in order to the admission of air, to the influence of which they should be gradually hardened.

SYMPHYTUM BOHEMICUM.

RED-FLOWERED COMFREY.

Class.
PENTANDRIA.

Order.
MONOGYNIA.

Natural Order.
BORAGINÆ.

Native of Bohemia.	Height. 8 inches.	Flowers in May, June.	Duration. Perennial.	Cultivated in 1801.
-----------------------	----------------------	--------------------------	-------------------------	------------------------

No. 71.

A Greek name corresponding with this was in use amongst the ancients, and was, by Tournefort, referred to this genus. Its signification was, to cement, or conglutinate, which was thought applicable to these plants, either from their glutinous juices, or from their healing qualities. Bohemicum, from Bohemia, its native country.

This species may probably be considered a mere variety of the *Symphytum officinale*. Such distinctive characters as it possesses appear, however, to be permanent.

The very powerful virtues of the Comfrey are well recorded by Camerarius, who saith that, "The rootes being outwardly applyed, helpeth fresh wounds or cuts immediately; being bruised and laid thereto, by glueing together their lips, and is especiall good for ruptures and broken bones; yea it is said to be so powerfull to consolidate or knit together, whatsoever needeth knitting, that if they be boyled with dissevered pieces of flesh in a pot, it will joyne them together againe."

It is easily propagated by dividing the roots, or by seeds; and will grow in any common garden soil.

Schmidt Bohem. n. 211.

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

THE JOURNAL OF THE ROYAL ANTHROPOLOGICAL INSTITUTE
Vol. 11, Part 1, 1881

DAPHNE GNIDIUM.

FLAX-LEAVED DAPHNE.

Class.
OCTANDRIA.

Order.
DIGYNIA.

Natural Order.
THYMELÆÆ.

Native of Spain.	Height. 2 feet.	Flowers in June, July.	Duration. Perennial.	Cultivated in 1597.
---------------------	--------------------	---------------------------	-------------------------	------------------------

No. 72.

As some of this genus bear a resemblance to the laurel, Linneus distinguished it by the name *Daphne*, in allusion to the fabled transformation of that nymph into such shrub. This species of *Daphne* retains the trivial name *Gnidium*, from the probability of its being the true GNIDION of the Greeks; who named it after *Gnidus*, a promontory and town in Asia Minor, where Venus had her temple.

Though the subjects of the genus *Daphne* are principally natives of Europe, yet, in their general habits, they class with that beautiful tribe, the evergreen American plants. Many of the most beautiful of these American shrubs are the under-wood of the extensive woodlands possessed by that quarter of our globe. In such situations, shaded in the summer from the mid-day sun, and in the winter, screened from cutting winds, and further protected and manured by the fallen leaves of the forest trees, they grow with a luxuriance that we can scarcely hope to witness in our gardens.

The *Daphne Gnidium* is chiefly propagated by being grafted on the *Daphne laureola*, or Spurge Laurel, and should be planted in sandy peat.

Hort. Kew. 2, v. 2, 410.

DAPHNE GENKWA

FLAX-LEAVED DAPHNE

INDIA
NORTH

CHINA
SICHUAN

General Description
HYPERICACEAE

Native or Sp. No.	Height in feet	Flower in June, July	Location General	Cultivated in 1907
----------------------	-------------------	-------------------------	---------------------	-----------------------

No. 72

As some of the genus bear a resemblance to the laurel, I named this distinguished it by the name *Daphne*, in allusion to the fabled transformation of that nymph into such shrub. This species of *Daphne* retains the trivial name *Genkwa*, from the probability of its being the true *Genkwa* of the Chinese; who named it after *Genkwa*, a promontory and town in Asia Minor, where 7 canals had her temple.

Though the subjects of the genus *Daphne* are principally natives of Europe, yet, in their general habits, they class with that beautiful tribe, the evergreen American plants. Many of the most beautiful of these American shrubs are the under-wood of the extensive woodlands possessed by that quarter of our globe. In such situations, shaded in the summer from the mid-day sun, and in the winter, screened from cutting winds, and further protected and manured by the fallen leaves of the forest trees, they grow with a luxuriance that we can scarcely hope to witness in our gardens.

The *Daphne Genkwa* is chiefly propagated by being grafted on the *Daphne laurifolia*, or *laurifolia*, and should be planted in sandy soil.

Herb. Kew. 9. v. 2. 410





Lobelia fulgens.

$\frac{1}{2}$



Erica Mediterranea.



Gaurn biennis.



Cacalia coccinea.

$\frac{2}{3}$

LOBE'LIA FUL'GENS.

FULGENT LOBELIA.

Class.

PENTANDRIA.

Order.

MONOGYNIA.

Natural Order.

CAMPANULACEÆ.

Native of	Height.	Flowers in	Duration.	Introduced
Mexico.	4 feet.	June, Sep.	Perennial.	in 1809.

No. 73.

This genus was named after Matthias de Lobel, see No. 61. Fulgens, from the Latin, bright, very conspicuous.

It may be conceived that every individual who feels an interest in the beauty of flowers, and in the possession of plants, bearing evidence, by their luxuriance, of superior cultivation, will be highly gratified in being informed how this magnificent herbaceous Lobelia may be produced in the highest possible perfection and splendour.

Mr. Sabine observes that 'it has lately been found to bear the severity of our winter, by being immersed in water, as an aquatic; and with this treatment has flowered well by the sides of ponds and in cisterns; but it was reserved for the skill of Mr. W. Hedges to discover a mode of culture under which this beautiful exotic has assumed a character of magnificence which will hereafter make it one of the most conspicuous decorations of our flower gardens.'

Mr. Hedges, in a communication which is printed in the second volume of the Horticultural Society's Transactions, directs the offsets to be divided in October, put into small pots, protected by a cold frame

till the middle of January. To be then removed to a hotbed or pine pit, and repotted, at intervals, till May; then to be taken into a greenhouse till they begin to flower.

We recommend the same principles, but vary the application of them a little, that they may be more generally useful. The offsets need not be divided till the latter end of February; and then they should be planted singly in pots of rich soil, rendered very light by the addition of decayed leaves or other vegetable mould, with a good portion of sand, and be kept in a moderate hotbed, where plenty of air can be admitted during the day time. About the end of March, remove the plants, with the roots and soil complete, into pots a little larger than those first employed, filling up the space with the same compost as before. After this the transplanting should be repeated every six weeks, still using pots a little larger at each removal, till through a gradation of four or five sizes, from small ones of four inches, you arrive at those of not less than ten inches diameter. Sink the plants nearly an inch in each fresh pot, and observe to keep them, from the commencement, in pans, which should never be without water, as much of the success depends on their continual moisture.

They may be taken out of the hotbed about the end of May, or even earlier, provided a temporary covering be afforded them at night.

When the plants have done flowering, cut off the stems; and during the severity of winter protect the offsets in a cold frame or airy room, where they may remain with moderate waterings till they are required to be again divided.

ERICA MEDITERRA'NEA.

MEDITERRANEAN HEATH.

Class.
OCTANDRIA.

Order.
MONOGYNIA.

Natural Order.
ERICÆ.

Native of Portugal.	Height. 2 feet.	Flowers in April, May.	Duration. Perennial.	Cultivated in 1648.
------------------------	--------------------	---------------------------	-------------------------	------------------------

No. 74.

The name of this beautiful genus has been noticed under No. 22, as derived from the Greeks; and it is remarked by Dr. Sibthorp that a corruption of the term EREICO is still used by them, and applied to the several species of this genus. Mediterranea has, probably, been chosen as indicative of the inland situations of which this species is native.

This, like most others of the beautiful tribe to which it belongs, should be planted in sandy peat, and may be increased either by cuttings or layers. All the heaths should have occasional waterings in the heat of summer, or they may fall into excessive langour, as Stillingfleet observes of singing birds, after midsummer. This sultry season further reminds us of his observations on the same subject;—No longer stimulated by the enchantments of spring, and the growing love of their mates, birds fall into supinity and the indolence of age.

“The groves, the fields, the meadows, now no more
With melody resound. 'Tis silence all,
As if the lovely songsters, overwhelm'd
By bounteous nature's plenty, lay entranc'd
In drowsy lethargy.”

Hort. Kew. 2, v. 2, 367.

GAU'RA BIEN'NIS.

BIENNIAL GAURA.

Class.
OCTANDRIA.

Order.
MONOGYNIA.

Natural Order.
ONAGRARIÆ.

Native of N. America.	Height. 6 feet.	Flowers in Aug. Oct.	Duration. Biennial.	Introduced in 1762.
--------------------------	--------------------	-------------------------	------------------------	------------------------

No. 75.

The term Gaura has been deduced from the Greek GAUROS, signifying pompous or stately. Biennis, from its duration being of two years only.

This stately herbaceous plant has very properly been named Gaura, from its free and lofty growth, its luxuriant branches, and its display of showy flowers in the evening. Though each corolla that expands in the afternoon, closes on the following morning, yet its gaiety is maintained by the continued extension of its flowering stems, and the production of numerous young branches, which also yield their proportion of blossoms.

Seeds of the Gaura biennis may be sown in the autumn as soon as ripe, or at the latter end of February. The young plants should be kept thin and free from weeds during the summer; and in the autumn be carefully transplanted where they are to remain. In the following summer they will produce their lofty flowering stems, which should be properly confined to strong upright supporters, in order that they may be effectually protected against the rude winds that frequently occur in September and despoil their beauty.

Hort. Kew. 2, v. 2, 344.

GARDA BIDENTIS.

BIDENTIS GARDA.

Flower.
HONGKONG.

Flower.
HONGKONG.

Natural Order.
GRAMINEAE.

N. America.	Height. in feet.	Flowers in Aug. Oct.	Duration in months.	Introduced in 1768.
-------------	---------------------	-------------------------	------------------------	------------------------

No. 75.

The name *Garda* has been deduced from the Greek *GAURO*, signifying pompous or stately. *Bidentis*, from its duration being of two years only.

This stately herbaceous plant has very properly been named *Garda*, from its free and lofty growth, its luxuriant branches, and its display of showy flowers in the evening. Though each corolla that expands in the afternoon, closes on the following morning, yet its grainy is maintained by the continual extension of its flowering stems, and the production of numerous young branches, which also yield their proportion of blossoms.

Seeds of the *Garda bidentis* may be sown in the autumn as soon as ripe, or at the latter end of February. The young plants should be kept thin and free from weeds during the summer; and in the autumn be carefully transplanted where they are to remain. In the following summer they will produce their lofty flowering stems, which should be properly confined to strong upright supports, in order that they may be effectually protected against the rude winds that frequently occur in September and despoil their beauty.

Herb. Kew. 2. t. 252.

CACALIA COCCINEA.

SCARLET-FLOWERED CACALIA.

Class.
SYNGENESIA.

Order.
POLYGAMIA ÆQUALIS.

Natural Order.
CORYMBIFERÆ.

Native of	Height.	Flowers in	Duration.	Introduced
Uncertain.	18 inches.	Aug. Sep.	Annual.	in 1800.

No. 76.

The generic name we have now to notice, was in use by Dioscorides, a celebrated Greek physician and botanist, who lived soon after the christian era. It is compounded of the two Greek words *CACON*, bad, and *LIAN*, exceedingly; from the real or supposed mischievous properties of the plant which bore the name, to the soil on which it grew. *Coccinea* from the Latin, scarlet or crimson-coloured.

According to Curtis, seeds of this plant were brought to England from Paris, in 1800; but to what country it is indigenous we are not correctly informed. It is a brilliant appendage to the parterre, in September; and contrasts well with the prevailing colours of that season.

This annual is of rather delicate habits, and the seed should be sown on a hotbed in the spring. Or they may be sown in pots, and put into a cucumber frame, where the young plants should not be crowded, but have as much air as can conveniently be allowed them. They may be planted into the open ground in the latter part of May, or at the beginning of June; and care should be taken that the roots be disturbed as little as possible.

Bot. Mag. 564.

CACALIA COCCINEA.

SCARLET-FLOWERED CACALIA.

Class. SYDNEY.
Order. POLYANTHA.
Natural Order. COMPOSITAE.

Native of	Height.	Flowers in	Duration.	Introduced
East India.	18 inches.	Aug. Sep.	Annual.	in 1800.


No. 78.

The generic name we have now in notice, was in use by Dioscorides, a celebrated Greek physician and botanist, who lived soon after the Christian era. It is compounded of the two Greek words *caca*, bad, and *lian*, exceedingly; from the real or supposed mischievous properties of the plant which bore the name, to the soil on which it grew. *Coc* comes from the Latin, scarlet or crimson-colored.

According to Curtis, seeds of this plant were brought to England from Paris, in 1800; but in what country it is indigenous we are not exactly informed. It is a brilliant appendage to the parterre, in September; and contrasts well with the prevailing colours of that season.

This annual is of rather delicate habit, and the seed should be sown on a hotbed in the spring. Or they may be sown in pots, and put into a greenhouse, where the young plants should not be crowded, but have as much air as can conveniently be allowed them. They may be planted into the open ground in the latter part of May, or at the beginning of June; and care should be taken that the roots be disturbed as little as possible.

Bot. Mag. 201.





Hibiscus Syriacus.

2



Asclepias tuberosa.



Oenothera purpurea.

2



Teucrium pyrenaicum.

HIBIS'CUS SYRIA'CUS.

ALTHÆA FRUTEX.

Class.

MONADELPHIA.

Order.

POLYANDRIA.

Natural Order.

MALVACEÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
Syria.	6 feet.	Aug. Sep.	Perennial.	in 1596.

No. 77.

Hibiscus is a name which has been handed down to us from the old Greek writers, but from what root the word originally sprung, is now unknown. As it was supposed to have been formerly applied to some of the mallow tribe, Linneus made choice of it to distinguish a splendid genus of the malvaceous order, of which our present specimen furnishes a good example. Syriacus, from Syria, which is its native country.

Our drawing of this species was taken from a beautiful variety, known by the appellation of the Painted Lady. There are, however, others, as the purple-flowered, stripe-flowered, white-flowered, double-flowered, variegated-leaved, &c.

Though varieties of the *Althæa frutex* have been cultivated in England during a space of 200 years, still we find this species by no means common. We have indeed been surprised to observe the scarcity of this beautiful shrub in plantations of no ordinary merit or extent; and we can attribute it only to partial failures in cultivation.

It has been propagated from seeds, cuttings, and layers; but so superior have seedling plants always

proved, that we shall only endeavour to supply directions for that mode of increase. Thus raised, they assume a more healthy habit, grow larger, and yield a display of much finer flowers. As the propagation of the *Hibiscus Syriacus*, from seeds, has not come fully under our immediate observation, we shall take the liberty of supplying the necessary information from what may be considered a good authority, Miller's Dictionary.

The seeds should be sown in pots, filled with light earth, about the end of March, and if they are placed in a gentle hotbed, it will greatly forward the growth of the young plants. When they are come up, they must be inured to the open air, and in May the pots should be plunged into the ground, in a border exposed to the east, where they may have the morning sun. By thus plunging the pots, the soil in them is prevented from drying so quickly as it would if they were left on the surface, and less attention is required in watering them during the summer. The plants should be kept free from weeds and tolerably moist; and in autumn it will be proper to remove the pots into a common frame, to screen them from frost; or into some other well-protected situation; for although these plants, when they have obtained strength, will resist the cold of our winters, yet the young plants, whose shoots are tender, are very often injured by the early frosts of autumn.

In the following spring they should be planted nine inches apart, in beds of light rich earth; be kept free from weeds: and in the winter again protected. Here they may remain one or two years, and should then be finally transplanted.

Hort. Kew. 2, v. 4, 226.

ASCLEPIAS TUBEROSA.

TUBEROUS SWALLOW-WORT.

Class.

PENTANDRIA.

Order.

DIGYNIA.

Natural Order.

ASCLEPIADEÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
N. America.	18 inches.	July, Aug.	Perennial.	in 1690.

No. 78.

Some portion of the plants comprised in this genus, must, no doubt, have been well known to the founder of our botanical system as possessing eminent medical qualities, or he would not have distinguished it by a name derived immediately from the god of medicine—Æsculapius. Its trivial name, tuberosa, may be applicable, in its more luxuriant state in America, but with us its roots are more fusiform than tuberous.

Its flowers are both singular and interesting; and where a suitable soil occurs for the growth of the plant, it should form a portion of every collection.

In America they call it the butterfly-weed, or pleurisy-root; and its medicinal qualities are highly appreciated. The root, when dry, is brittle and easily reduced to powder; and its taste is moderately bitter, but not otherwise unpleasant.

Dr. Bigelow says that it is eminently entitled to the attention of physicians, as an expectorant and diaphoretic. It produces effects of this kind with great gentleness, and without the heating tendency which accompanies many vegetable sudorifics. It appears to be an expectorant peculiarly suited to the

advanced stages of pulmonary inflammation, after depletion has been carried to the requisite extent.

Dr. Parker, of Virginia, having been in the habit of employing this root for twenty-five years, considers it as possessing a peculiar and almost specific quality of acting upon the organs of respiration; promoting suppressed expectoration, and relieving the breathing of pleuritic patients in the most advanced stages of the disease.

Like other vegetable bitters, if given in large quantities, especially in infusion and decoction, it operates on the alimentary canal, though its efficacy in this respect is not sufficient to entitle it to rank amongst active cathartics.

The best mode of administering the *Asclepias* root, is in decoction or substance. A teacup-full of the strong decoction, or from twenty to thirty grains of the powder, may be given in pulmonary complaints several times a day.

Success does not always attend the best efforts to preserve this plant. This generally arises from one of two causes;—that of removing old plants, or depositing them in moist situations. In America it is found in dry sandy soils, and pine woods; and attempts to preserve it in wet or stiff earth will generally prove abortive.

It should be raised from seeds, which, as they are not frequently perfected in England, must be obtained from America. These may be planted in spring, on a bed of light sandy earth, and it will be an advantage if they can be raised in the situations in which they are to remain. If transplanting be required, perform it when the plants are one year old.

Hort. Kew. 2, v. 2, 82.

ÆNOTHERA PURPU'REA.

PURPLE-FLOWERED ÆNOTHERA.

Class.
OCTANDRIA.

Order.
MONOGYNIA.

Natural Order.
ONAGRARIÆ.

Native of N.America.	Height. 18 inches	Flowers in July, Aug.	Duration. Annual.	Introduced in 1791.
-------------------------	----------------------	--------------------------	----------------------	------------------------

No. 79.

The derivation of the term Ænothéra, from the Greek, has been noticed under No. 35. Purpurea, purple, indicative of the colour of the flower.

This herbaceous plant, forms a pretty contrast, both in flower and foliage, to the usual variety of annuals which furnish our borders and mounts at its own period of flowering.

It is of moderate growth, never rising into extreme luxuriance, to the destruction of other subjects near to it, and generally supports itself without assistance.

The greater part of this genus expand their flowers in the evening, and their beauty fades on being exposed to the rays of the sun next morning. The present species, however, possesses the advantage of supporting its expanded flowers through the whole of the day.

Though frequently considered as a tender annual, the Ænothéra purpurea usually succeeds best when sown in light rich soil where it is to flower. Or it may be sown on a seed bed in March or April, and transplanted into the flowering compartment when the plants are two or three inches high.

Hort. Kew. 2, v. 2, 344.

TEU'CRIMUM PYRENA'ICUM.

PYRENEAN TEUCRIUM.

Class.
DIDYNAMIA.

Order.
GYMOSPERMIA.

Natural Order.
LABIATEÆ.

Native of Pyrenees.	Height. 3 inches.	Flowers in June, Aug.	Duration. Perennial.	Introduced in 1731.
------------------------	----------------------	--------------------------	-------------------------	------------------------

No. 80.

It is supposed that the word Teucrium has arisen out of the name of the Trojan Prince Teucer, the father-in-law of Dardanus, king of Troy, but on what account does not appear. As the country of Troy is sometimes called Teucra the name may originally have been given to some plant indigenous there. Pyrenaicum, from Pyrenees, where it is found. The English appellation, Germander, is rarely applied to any, excepting two or three British species; which have been, by some authors, regarded as possessing medicinal virtues.

The common Germander, or Teucrium Chamædrys, has been esteemed as beneficial in gout and rheumatism, and is one of the vegetables that constitute the celebrated Portland gout powder.

The Teucrium pyrenaicum, from its humble growth, is well suited to the fronts of borders, and for decorating artificial rock-work, where it will be found to grow in perfection.

It may be planted in any common garden soil, and increased by a division of its roots in spring; which time is preferable to autumn, as the plants make good roots before winter.

Hort. Kew. 2, v. 3, 371.

THE CRUIC PYRENEAICUM.

PYRENEAN TREES.

Order.
PYRENEAIC.

Class.
DIDYMARIA.

Natural Order.
LABIATAE.

Native of Pyrenees.	Height 3 feet.	Flowers in racemes, Aug. to Sept.	Latitudes 45° to 50°.
------------------------	-------------------	--------------------------------------	--------------------------

It is

It is supposed that the word *Toum* has arisen out of the name of the Trojan Prince *Troas*, the father-in-law of *Paris*, king of *Troy*, but no what account does not appear. As the country of *Troy* is sometimes called *Troas* the name may originally have been given to some plant indigenous there. *Pyreneum*, from *Pyrene*, where it is found. The English appellation, *Pyreneum*, is rarely applied to any, excepting two or three British species; which have been, by some authors, regarded as possessing medicinal virtues.

The common *Pyreneum*, or *Toum*, *Chamaedryas*, has been introduced as medicinal in goat and rheumatism, and is one of the vegetables that constitute the celebrated *Pyreneum* goat powder.

The *Toum* *pyreneum*, from its humble growth, is well suited to the fronts of houses, and for decorating artificial rock-work, where it will be found to grow in perfection.

It may be planted in any common garden soil, and increased by a division of its roots in spring; which time is preferable to autumn, as the plants make good roots before winter.

Bot. Rev. 2. 2. 271





Chelone barbata.

½



Iberis sempervirens.

½



Ramonda pyrenaica.

½



Vesicaria utriculata.

½

CHELO'NE BARBATA.

SCARLET CHELONE.

Class.
DIDYNAMIA.

Order.
ANGIOSPERMIA.

Natural Order.
BIGNONIACEÆ.

Native of Mexico.	Height. 4 feet.	Flowers in July, Aug.	Duration. Perennial.	Introduced in 1794.
----------------------	--------------------	--------------------------	-------------------------	------------------------

No. 81.

The term Chelone, is derived from a similar word in the Greek language, signifying a tortoise. Barbata, bearded.

This interesting herbaceous plant, with its delicate slender stems, supporting a multitude of beautiful pendent scarlet flowers, is surpassed in elegance but by few subjects of similar magnitude. It is never intrusive by its foliage or stems, and simply requires the support of a thin willow shoot, as a guard against occasional winds. It is the Chelone Rueillioides of Andrews's Repository.

It should be planted in a dry warm situation and southerly aspect; and in severe frosts should be covered with a hand-glass or coarse straw. It may be propagated by occasional divisions of the root, which is best effected late in the spring.

A certain method to prevent disappointment, is to take cuttings of the young shoots as early as they will admit of it. These may be planted under a hand-glass, and should be potted after they are well rooted. When frosts set in, give them the protection of the cold frame, and in April they may be turned into the borders.

CHLOEON BARBATA.

SECRET CHLOEON

Native of	Height	Flowers in
Mexico	4 feet	July, Aug. to 1792.

Dec. 21.

The term *Chloëon*, is derived from a similar word in the Greek language, signifying a testicle. *Barbata*, Latin, bearded.

This is a very herbaceous plant, with its delicate slender stems, supporting a multitude of beautiful pendant scarlet flowers, is surpassed in elegance by few subjects of similar magnitude. It is never intruded by its foliage or stems, and simply requires the support of a thin wither shoot, as a guard against occasional winds. It is the *Chloëon* *Humboldtii* of Andrews's Repository.

It should be planted in a dry warm situation and southerly aspect; and in severe frosts should be covered with a hand-glass or warm glass. It is propagated by occasional divisions of the root, which is best selected late in the autumn.

A certain method to prevent the plant from taking entrance of the young shoots as early as the will admit of it. These may be placed in a warm sand glass and should be potted after the manner of the *Chloëon* *Humboldtii*, and in April or May, when the young shoots are two feet high.

1802. Nov. 2. 1. 1. 1.

IBERIS SEMPERVIRENS.

NARROW-LEAVED CANDY-TUFT.

Class.
TETRADYNAMIA.

Order.
SILICULOSA.

Natural Order.
CRUCIFERÆ.

Native of Candia.	Height. 9 inches.	Flowers in April, May.	Duration. Perennial.	Cultivated in 1731.
----------------------	----------------------	---------------------------	-------------------------	------------------------

No. 82.

The name of this genus is one retained from Dioscorides; and was applied by him to some plant resembling the present one. The term is supposed to have been originally deduced from Iberia, a name used by the Greeks for Spain; where, possibly, the Iberis of the Greeks may have been first noticed. Sempervirens is an appellation compounded from the Latin, signifying always green, in allusion to its evergreen habits.

It is a most desirable little shrub; for as well as decorating the garden with its beautiful white tufts of flowers, during two months of the spring, it exhibits, in the winter, by its evergreen foliage, a lively remembrance of the verdure that is past, and also a foretaste of that which we are happy to anticipate as again to come. Though winter may occasionally seem to conquer its tenacity for life, yet, no sooner does the severest frost relax its icy grasp, than the Iberis sempervirens appears again in spring-like freshness, to exult in its regained liberty.

It is very readily propagated, either by fastening down its branches beneath the soil, or by cuttings taken in the spring.

Hort. Kew. 2, v. 4, 83.

IBERIA SEMPERVIRENS.
NARROW-LEAVED CANDY-TREE.

Class. TETRADYMIA.
Order. Natural Order.
CROZIERIA.

Native of	Height.	Flowers in	Position.	Collected
Florida.	9 inches.	April, May.	Vertical.	in 1734.

No. 82.

The name of this genus is one retained from Di-
scorides; and was applied by him to some plant
resembling the present one. The term is supposed
to have been originally deduced from Iberia, a name
used by the Greeks for Spain; where, possibly, the
Iberis of the Greeks may have been first noticed.
Sempervirens is an appellation transferred from
the Latin, signifying always green, in allusion to its
evergreen habit.

It is a most desirable little shrub; for as well as
decorating the garden with its beautiful white tufts
of flowers, during two months of the spring, it exhi-
bits, in the winter, by its evergreen foliage, a lively
remembrance of the verdure that is past, and also a
foretaste of that which we are happy to anticipate
as again to come. Though winter may occasionally
seem to deprive its beauty for life, yet no sooner
does the severest frost relax its icy grasp, than the
Iberis sempervirens appears again in spring-like
freshness, to exult in its renewed liberty.

It is very readily propagated, either by inserting
down its branches beneath the soil, or by cuttings
taken in the spring.

Flora, &c. &c. A. 82.

RAMONDA PYRENAICA.

BORAGE-LEAVED RAMONDA.

Class.

PENTANDRIA.

Order.

MONOGYNIA.

Natural Order.

SOLANEE.

Native of Pyrenees.	Height. 5 inches.	Flowers in May, June.	Duration. Perennial.	Cultivated in 1731.
------------------------	----------------------	--------------------------	-------------------------	------------------------

No. 83.

This genus received its title in honour of a French botanist, of the name of Ramond. *Pyrenaica*, from Pyrenees, where it is indigenous.

The present beautiful alpine plant is the *Verbascum myconi* of Linneus, Curtis, and Aiton, and consequently is now commonly known by that name. Our aversion to unnecessary innovations on the established arrangement and nomenclature of botanical science has been fully expressed; and it is only from conviction in our own humble judgment, of the necessity or expediency of alteration, that we ever submit to changes. As it is, in all cases, necessary on the one hand, to guard against the intrusion of empirics; so, on the other, it is expedient that we attach not ourselves, by undue prejudice, to any system of things, merely on account of a long acquaintance with it.

Brilliant genius sometimes steps forth and strikes out a new and enlightened path for itself, but unfortunately, a great portion of the innovators on all received systems, are found to pursue a road, ultimately deserted by all but themselves. An anonymous author observes, that a virtuous mind has

primarily a sense of justice, which teaches a regard to the rights of others, among which rights are their opinions.

The *Ramonda Pyrenaica* is a desirable little plant to cultivate, either in pots or in the borders; and should occupy a place amongst the various low close-growing alpine subjects. It may be increased by dividing the roots, or by seeds; and succeeds best in a cool situation.

It is also a suitable subject for pot culture, and we cannot give our readers more judicious advice, respecting its management, than is contained in the remarks of a correspondent; who says, that the various trials that I have made with the *Ramonda Pyrenaica*, have satisfied me that bog-earth is better adapted to its habits of growth than a more substantial and retentive soil. Indeed, I have long made it a rule to provide plants with food, rather with reference to the fibres of their roots, than to their apparent wants of strong or mild nourishment. All plants which have wiry roots, I invariably find, delight in a peat mould, with a good proportion of the decayed roots of the peat, and a little white sand mixed up with it; and if planted in pots, well drained, first with very small broken tiles, and next with nothing but decayed roots of peat earth, they invariably thrive. This plan enables the broken tiles to answer their intended end much longer than they could possibly do, were they placed immediately in contact with the finely sifted mould; and upon examining the roots of a plant so treated, you will find the fibres spread over, and freely intermixed with, the drainers.

Hort. Kew. 2, v. 1, 386.

VESICARIA UTRICULATA.

SMOOTH VESICARIA.

Class.
TETRADYNAMIA.

Order.
SILIKUOSA.

Natural Order.
CRUCIFERÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
Levant.	13 inches.	April, June.	Perennial.	in 1739.

No. 84.

Vesicaria, from the Latin vesica, a bladder, in allusion to its bladder-like seed vessels. Utriculata is also from the Latin, and has nearly the same signification.

This plant is the *Alyssum utriculatum* of Curtis and others ; but from its appearance in the *Hortus Kewensis*, under the genus *Vesicaria*, it has thence been copied into the general catalogues, and is now distinguished in most respectable nurseries, by the name we have adopted.

It is an extremely gay and hardy herbaceous plant, continuing in bloom a considerable time. After its blossoms have faded, its spike of inflated silicles still form an object by no means uninteresting, till the seeds are ripe in July. It will flower in any common garden soil, and may be readily propagated either by seeds or from cuttings. Probably the following method of increase may be pursued with advantage. Fill up the interstices of the plant in the summer with soil, so that the whole of the branches may, in reality, be laid ; they will, it is presumed, make strong plants for separation in the following year.

Hort. Kew. 2, v. 4, 97.

VERICARIA UTRICULATA.

SMOOTH VERICARIA.

Vericaria utriculata is a biennial or triennial plant, with a creeping rootstock, and a very slender, upright stem, which is branched at the top, and is furnished with small, opposite, ovate leaves, which are very smooth, and have a very short petiole. The flowers are small, and are produced in a loose, terminal raceme. The fruit is a small, round, bladder-like capsule, which is attached to the stem by a long, slender pedicel.

Native of	Europe, Asia, and Africa.
Height	1 to 2 feet.
Flowers in	June, July, and August.
Fructification	in July.

This plant is the African vericaria of Clinton and others; but from its appearance in the Italian Peninsula, under the name Vericaria, it has been distinguished in most respectable nurseries, by the name we have adopted. It is an extremely gay and hardy herbaceous plant, containing no blood, a considerable quantity of which is contained in its roots, and is often used in the East for medicinal purposes. It will flower in any common garden soil, and may be readily propagated either by seeds or from cuttings. Probably the following method of increase may be pursued with advantage. Fill up the intervals of the plant in the summer with soil, so that the whole of the branches may, in winter, be buried; they will, in the following year, produce strong plants for propagation in the following year.

Herb. Acad. 1757, p. 57.





Argemone Mexicana.

1/2



Statice oleaeifolia.



Lychnis chalcedonica.

1/2



Anthericum liliastrum.

ARGEMONE MEXICANA.

MEXICAN ARGEMONE.

Class.

POLYANDRIA.

Order.

MONOGYNIA.

Natural Order.

PAPAVERACEÆ.

Native of America.	Height. 2 feet.	Flowers in July, Aug.	Duration. Perennial.	Cultivated in 1597.
-----------------------	--------------------	--------------------------	-------------------------	------------------------

No. 85.

The name of this genus, Argemone, has been handed down to us from the ancient Greek botanists. They adopted it after the term ARGEMA, a disease of the eyes; wherein white spots arise on the cornea, which their argemone was said to be efficacious in removing.

Gerard says, 'The golden Thistle of Peru, called in the West Indies, Figue del Inferno, (Infernal Fig,) a friend of mine brought it unto me from an island there, called Saint John's Island, among other seeds. What reason the inhabitants there have to call it so, is unto me unknown, unless it be because of his fruit, which doth much resemble a fig in shape and bigness, but so full of sharp and venomous prickles, that whosoever had one of them in his throat, doubtless it would send him packing either to heaven or hell.'

Those of our readers who happen to possess Johnson's edition of Gerard's Herbal, should be informed, that the principal description of the above plant is contained in that work, at page 1155; but the wood-cut figure of it is at page 371, and is mis-numbered 401.

It is said to be common in Mexico, and all the islands of the West Indies, where it is a troublesome weed in their cultivated lands, and has been found in a wild state in some of the southern countries of Europe.

The inspissated juice of the leaves and stems, forms a pigment, in colour between sap green and gamboge, but apparently not more valuable to the artist than a mixture of those substances. It is esteemed very detersive, and is generally used in diseases of the eyes; but the infusion is looked upon as a sudorific and resolute, which may be used with success on many occasions.

The seeds are said to be a much stronger narcotic than opium. They are thought to be an excellent remedy, and are frequently administered by the inhabitants, in the sugar colonies, in diarrhoea and the bloody flux. They have a trifling degree of pungency, but it does not manifest itself for some time upon the palate.

The exterior covering of the seed assumes the appearance of delicate net-work, which becomes more marked and prominent, as they ripen and dry; and the hilum, or eye, forms a fine seam on one side of it, similar to an artificial enclosure.

There is no difficulty in cultivating the *Argemone Mexicana*, as an annual; excepting, that like many others of the papaveraceous tribe, they are impatient of removal. It is better to sow them in a light soil, in the spring, where they are to remain; and if the seed, when ripe, be scattered from the plants, they will generally vegetate in the spring without further trouble.

STATICE OLEÆFO'LIA.

OLIVE-LEAVED SEA LAVENDER.

Class.
PENTANDRIA.

Order.
PENTAGYNIA.

Natural Order.
PLUMBAGINEÆ.

Native of Italy.	Height. 15 inches.	Flowers in July, Aug.	Duration. Perennial.	Cultivated in 1683.
---------------------	-----------------------	--------------------------	-------------------------	------------------------

No. 86.

Much uncertainty exists respecting the original signification of the generic term *Statice*. It is supposed to have arisen out of the Greek *STATIZO*, to stop or arrest, in allusion to its astringent quality. Sir J. E. Smith observes, that what the ancient plant may have been, can scarcely be guessed with any probability. The modern application of the name to our Thrift or Sea-Gilliflower, he observes, seems to have originated with Dalechamp, whom Tournefort followed. Hence it has become appropriated to a fine and extensive genus, whose wiry and entangled stems, so well formed to impede the progress of a foot passenger, may literally almost justify its present use.

The present species is smaller than the greater portion of them, but still is an interesting and perfectly hardy little plant. Our figure of a blossom-branch is the full size, but the radical leaves are only half the size of nature.

It flourishes in a light loam, and may be increased by a division of the roots. It does not increase very fast; and will always flower stronger from having remained two years without a removal.

Hort. Kew. 2, v. 2, 181.

LYCHNIS CHALCEDONICA. flore pleno.

DOUBLE SCARLET LYCHNIS.

Class.
DECANDRIA.

Order.
PENTAGYNIA.

Natural Order.
CARYOPHYLLÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
Russia.	3 feet.	June, July.	Perennial.	in 1596.

No. 87.

The origin of the name of this genus, like that of our last, is wrapped in uncertainty.

The Greeks used a similar name, which word also signifies a lamp. Hence conjectures arose, and ingenuity has been exerted, to trace the connexion. The term was formerly used for more plants than it now is; some of which may have admitted comparisons not applicable to the present *Lychnis* tribe. It is said that the down of the plant may have been used to make wicks; or that the colour of the flower was brilliant, as flame; also that the transparent membranous calyx resembled a lamp or lantern. *Chalcedonica*, from *Chalcedon*; whence seeds have been brought.

This splendid herbaceous plant, single or double, is highly ornamental, and should not be dispensed with. Parkinson, 200 years ago, notes it as a glorious flower, being then as rare as it is beautiful.

The single variety may be raised from seeds. The double or single may be increased by dividing the roots, or by cuttings of the stems, taken in June. It should be planted in a strong fresh loam, and have pure air.

Hort. Kew. 2, v. 3, 132.

LYCHNIS CHALCEDONICA. *Red plant.*
 DOUBLE SCARLET LYCHNIS.

Class. *Malvaceae.*
 Order. *Malvales.*
 Family. *Malvaceae.*
 Genus. *Lychnis.*
 Species. *Lychnis chalcedonica.*

Native of	Height	Flowers in	Flowers in	Flowers in
Asia.	3 feet	June, July.	Perennial.	is 1820.

No. 87.

The origin of the name of this genus, like that of our last, is wrapped in uncertainty. The Greeks used a similar name, which word also signifies a lamp. Hence conjecture arose, and ingenuity has been exerted, to trace the connection. The term was formerly used for more plants than it now is; some of which may have admitted comparison not applicable to the present *Lychnis* tribe. It is said that the down of the plant may have been used to make wicks; or that the color of the flower was brilliant, as flame; also that the transparent membranous cavity resembled a lamp or lantern. Chalcedonica, from Chalcedon; whence seeds have been brought.

This splendid herbaceous plant, single or double, is highly ornamental, and should not be dispensed with. Parkinson, 200 years ago, notes it as a glaucous flower, being then as rare as it is beautiful.

The single variety may be raised from seeds. The double or single may be increased by dividing the roots, or by cuttings of the stems, taken in June. It should be planted in a strong fresh loam, and have pure air.

ANTHERICUM LILIASTRUM.

SAVOY SPIDERWORT; OR, ST. BRUNO'S LILY.

Class.
HEXANDRIA.

Order.
MONOGYNIA.

Natural Order.
ASPHODELEÆ.

Native of Switzerland	Height. 18 inches.	Flowers in May, June.	Duration. Perennial.	Cultivated in 1629.
--------------------------	-----------------------	--------------------------	-------------------------	------------------------

No. 88.

Anthericum is deduced from two Greek words; ANTHOS, a flower; and REKOS, a hedge.

Liliastrum may signify star lily. The term has probably been transferred from some other plant to the present species. It was found wild in Savoy, and formerly called Phalangium, the name of a venomous species of spider, from its being considered an antidote to the bite of that insect. Hence comes our term Spiderwort. The French, we believe, inscribed it to St. Bruno, the celebrated founder of the Carthusians.

Linneus first considered it an Hemerocallis; but afterwards an Anthericum. Botanists are divided in opinion on this subject, and some now term it Hemerocallis liliastrum.

It increases but slowly, and if too often divided, will either not flower at all, or produce a diminutive show of blossoms. Autumn is the proper season for removing it, and if planted in a rather shady situation, though not immediately under the branches of shrubs or trees, it will succeed very well; and the duration of its delicate flowers, will be lengthened by shade from the direct rays of the sun.

Hort. Kew. 2, v. 2, 269.

ANTHRICUM LILASTRUM

SAVOY SPIRKEWORT; OR, ST. BRUNO'S LILY.

Order
MONOCOTYLEDONALES

Class
GRAMINEAE

Natural Order
ANTHRACEAE

Native of	Height	Flowers in	Duration	Cultivated
Subalpine 15 inches	15 inches	May, June	Perennial	in 1838


No. 87.

Anthriscum is deduced from two Greek words; *anthos*, a flower; and *arkos*, a hedge. *Lilastrum* may signify star lily. The term has probably been transferred from some other plant to the present species. It was found wild in Savoy, and formerly called *Phedragium*, the name of a venomous species of spider, from its being considered an antidote to the bite of that insect. Hence comes our term Spikewort. The French, we believe, inscribed it to St. Bruno, the celebrated founder of the Carthusians.

Linnæus first considered it an *Hemerocallis*; but afterwards an *Anthriscum*. Botanists are divided in opinion on this subject, and some now term it *Hemerocallis lilastrum*.

It increases but slowly, and is too often divided, will either not flower at all, or produce a diminutive show of blossoms. Autumn is the proper season for removing it, and it planted in a rather shady situation, though not immediately under the branches of shrubs or trees, it will succeed very well; and the duration of its delicate flowers, will be lengthened by shade from the direct rays of the sun.

Herb. Kew. 2. p. 200.





Menziesia globularis.



Cytisus capitatus

45



Chrysanthemum tricolor

42



Chelone obliqua

35

MENZIE'SIA GLOBULARIS.

GLOBE-FLOWERED MENZIESIA.

Class.
OCTANDRIA.

Order.
MONOGYNIA.

Natural Order.
RHODORACEÆ.

Native of N. America.	Height. 3 feet.	Flowers in June.	Duration. Perennial.	Introduced in 1806.
--------------------------	--------------------	---------------------	-------------------------	------------------------

No. 89.

This genus was named by Sir J. E. Smith, in honour of his friend Archibald Menzies; who made a voyage round the world with Vancouver, and collected many rare and unknown plants, particularly cryptogamic subjects. Globularis, from its globose corolla.

This deciduous little shrub cannot, for splendour, be ranked with the specious and imposing specimens of flowering subjects which generally constitute the natural order Rhodoraceæ, many of which have emanated from the same source as our present plant, North America, that rich fountain of vegetable beauty. It is a compact growing shrub, of rather slow growth, and frequently not exceeding twelve or eighteen inches in height. It should be planted in the foreground of the shrubbery, or American compartment; and its foliage will form a pleasing contrast with the deep glossy verdure of the Rhododendron, and evergreens of taller growth, which commonly give depth of shadow to the boundaries of cultivated grounds.

It should be planted in a mixture of peat and loam; and may be propagated by layers.

MEXICANA GLOBE-LIPS

ALBINO-FLOWERED MEXICANA

Height	Flowers in	Flowers in
1 foot	June	July

No. 10

This plant was named by Mr. J. E. Smith, in honor of his friend Archibald Menzies, who explored regions round the world with Vancouver, and collected many rare and unknown plants, particularly cryptogamic subjects. (Globe-lips, from its globe-like shape.)

This deciduous little shrub cannot, for aphyllous, be ranked with the evergreen and impenetrable specimens of flowering subjects which generally constitute the natural order Rhododendron, many of which have originated from the same source as our present plant. North America, that rich fountain of vegetable life. It is a compact growing shrub, of rather slow growth, and frequently and exceedingly sterile in culture unless in height. It should be placed in the background of the shrubbery, or American landscape; and its foliage will form a pleasing contrast with the deep green verdure of the Rhododendron, and (not) of taller growth, which commonly give depth of shadow to the landscape, or cultivated grounds.

It should be planted in a mixture of peat and sand, and may be propagated by layers.

From New York, N. Y.

CYTISUS CAPITATUS.

HEADED CYTISUS.

Class.
DIADELPHIA.

Order.
DECANDRIA.

Natural Order.
LEGUMINOSÆ.

Native of Austria.	Height. 2 feet.	Flowers in June, July.	Duration. Perennial.	Introduced in 1774.
-----------------------	--------------------	---------------------------	-------------------------	------------------------

No. 90.

A term similar to our *Cytisus* was in use amongst the ancient Greeks, whence ours has been derived. The particular plant which then bore the appellation was said, by Pliny, to have been found in the isle of Cythnus, one of the Cyclades, from which circumstance the Greek name is supposed to have originated. *Capitatus*, from the Latin, growing with a head; in allusion to its terminal mode of inflorescence, shown by the annexed representation.

The *Cytisus*, of which about twenty shrubby species are cultivated in England, is a general favourite in the pleasure grounds. The *Cytisus Laburnum* is universally known, and as generally admired; and our present subject, though far more humble, is by no means wanting in attraction.

This species of *Cytisus*, like several others, ripens its seed in our climate; and young plants are more conveniently raised from them than by any other means. They should be sown in a light soil, in March, and in about two months the young plants will appear. At one year old they should be transplanted to a nursery bed, and finally moved in the following spring.

Hort. Kew. 2, v, 4, 320.

CYTISUS CAPITATUS HEADED CYTISUS

CYTISUS
HEADED CYTISUS

CYTISUS
HEADED CYTISUS

HEADED CYTISUS
HEADED CYTISUS

Native of	Height	Flowers in	Duration	Perennial
Asia	2 feet	June, July	Perennial	in 1772

No. 66.

A form similar to our *Cytisus* was in use amongst the ancient Greeks, whence ours has been derived. The particular plant which then bore the appellation was said, by Pliney, to have been found in the island of Cyprus, one of the *Cyprides*, from which *Cyprum* stems the Greek name is supposed to have originated. *Capitatus*, from the Latin, growing with a head; in allusion to its terminal mode of inflorescence, shown by the annexed representation.

The *Cytisus*, of which about twenty shrubby species are cultivated in England, is a general favorite in the pleasure grounds. The *Cytisus Laburnum* is universally known, and is generally admired; and our present subject, though far more humble, is by no means wanting in attraction.

This species of *Cytisus*, like several others, ripens its seed in our climate; and young plants are more conveniently raised from them than by any other means. They should be sown in a light soil, in March, and in about two months the young plants will appear. At one year old they should be transplanted to a nursery bed, and finally moved in the following spring.

Seed, Nov. 2, 1772.

CHRYSAN'THEMUM TRI/COLOR.
THREE-COLOURED CHRYSANTHEMUM.

Class.
 SYNGENESIA.

Order.
 POLYGAMIA SUPERFLUA.

Natural Order.
 CORYMBIFERÆ.

Native of Barbary.	Height. 2 feet.	Flowers in July, Sep.	Duration. Annual.	Introduced in 1796.
-----------------------	--------------------	--------------------------	----------------------	------------------------

No. 91.

Chrysanthemum is derived from two Greek words, CHRUSOS, gold, and ANTHOS, a flower; and the brilliant golden hue of some species of Chrysanthemum fully justify the appellation.

In the variety of elegant annuals, which usually adorn our gardens, the Chrysanthemum tricolor should never be wanting. It flowers rather earlier than the common annual chrysanthemum, and the stems are not so branching nor obtrusive in their growth.

Having, so long, been accustomed to meet this plant under the appellation above given, we unconsciously wrote tricolor in directing the engraver. Chrysanthemum carinatum has of late been adopted; and it must be confessed there is a plausible reason for the exchange of nomenclature, when we are told that a variety of this species produces perfectly yellow flowers.

Seeds should be collected from such plants as grow at some distance from the Chrysanthemum coronarium, or common species; as we find them prone to mingle, to the injury of both. It may be sown with the common annuals.

Hort. Kew. 2, v. 5, 95.

CHRYSANTHEMUM TRICOLOR. THREE-COLORED CHRYSANTHEMUM.

Order. ASTROCARPÆA.
Suborder. POLYANTHA.
Tribe. POLYANTHÆ.
Genus. CHRYSANTHEMUM.

Native of	Height	Flower is	Duration	Introduced
Barbary	2 feet	July, Sep.	Annual	in 1790.

No. 91.

Chrysanthemum is derived from two Greek words, *chrysos*, gold, and *anthos*, a flower; and the brilliant golden hue of some species of *Chrysanthemum* fully justify the appellation.

In the variety of elegant annuals, which occasionally adorn our gardens, the *Chrysanthemum tricolor* should never be wanting. Its flowers rather earlier than the common annual *chrysanthemum*, and the stems are not so branching nor obtrusive in their growth.

Having, so long, been accustomed to meet this plant under the appellation above given, we unnecessarily wrote *tricolor* in directing the engraver. *Chrysanthemum carinatum* has of late been adopted; and it must be confessed there is a parallel reason for the exchange of nomenclature, when we are told that a variety of this species produces perfectly yellow flowers.

Seeds should be collected from such plants as grow at some distance from the *Chrysanthemum carinatum*, or common species; as we find them prone to adhere, to the injury of both. It may be sown with the common annuals.

Flora, &c. &c. 2. 2. 2.

CHELO'NE OBLI'QUA.

RED-FLOWERED CHELONE.

Class.

DIDYNAMIA.

Order.

ANGIOSPERMIA.

Natural Order.

BIGNONIACEÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
N.America.	2 feet.	Aug. Sep.	Perennial.	in 1752.

No. 92.

This genus was named by Tournefort after the Greek CHELONE, a tortoise; from a fancied resemblance of the flowers to that animal; its corolla being convex above and flat beneath. Obliqua, from the Latin, in allusion to the oblique position of its foliage.

This hardy herbaceous plant has long been a favourite amongst us, which may arise, not alone from its bold and handsome flowers, but from the little care it requires at our hands. Planted in any common soil that is tolerably retentive and moist, it is sure to succeed. We have occasionally seen it in a very light and dry border increase but little, and by its creeping roots change its situation, apparently in quest of nutriment, not afforded in sufficient quantity in its former residence. It has proceeded by its creeping roots, or rather it may be said, by its underground stems, to a distance of eighteen inches; there having halted, as it were, to colonize, like a parent directing his offspring, or a husbandman his labourers, has sent out its numerous fibres, to collect food from the surrounding soil. The fair author of an interesting little work, "The Wonders of the

Vegetable Kingdom," has very aptly compared the roots of a tree to the labouring classes of society. 'Society at large,' she observes, 'may be compared to a tree. The poor may be designated by the roots; the middle classes by the stem and branches; the dignified and noble, as well as those who adorn and improve humanity, by the flowers, leaves, and fruit. The stem is dependant on the root; without the stem the root would soon decay; flowers, fruits, and leaves, are equally ornamental and important to the parent tree. One member of the vegetable body cannot say unto another, I have no need of thee. To each an allotted duty is assigned; severed, they are of little worth; united, they form a beautiful and perfect whole.'

The various modes of reproduction, and the vegetable economy, exhibited in the growth and the adaption of the habits of roots, to the peculiar necessities of the plant, are well worthy the attention of every enquiring naturalist.

Some species of grass which, in moist situations, emit fibrous roots alone, will, in more uncongenial and dry ones, form small bulbs, whereby a reservoir of nutriment is secured against the occurrence of an irregular supply of the fluids requisite for its sustenance. Thus is shown to us the care of the Almighty over the smallest of his works.

The roots of large trees also, in unpropitious situations, have been observed to vary their natural mode of growth, most materially, in conformity with their need of nourishment. Some curious instances of such circumstances we may hereafter have occasion to notice.





Spigelia Marilandica.

35



Digitalis lutea.

36



Centaurea suaveolens.

37



Primula farinosa.

38

SPIGELIA MARILANDICA.

PERENNIAL WORM-GRASS, OR INDIAN PINK.

Class.

PENTANDRIA.

Order.

MONOGYNIA.

Natural Order.

GENTIANEÆ.

Native of N. America.	Height. 2 feet.	Flowers in June, July.	Duration. Perennial.	Introduced in 1694.
--------------------------	--------------------	---------------------------	-------------------------	------------------------

No. 93.

This genus was distinguished by Linneus, in commemoration of Adrian Spigelius, a botanist and physician of considerable acquirements, who was born at Brussels, in 1578. His "Isagogue in Rem Herbariam," published at Padua in 1606, is said to contain much interesting matter respecting the virtues of plants; and it is somewhat curious, that Spigelius collected a great portion of this knowledge from the peasants of Italy, by making a tour amongst them in the character of a rustic. Marilandica, from Maryland, one of the United States of America, where this plant is indigenous.

This is a beautifully ornamental herbaceous subject, that withstands the cold of our winters tolerably well, but does not increase much, and is not unfrequently lost. The dried stems and leaves are known to almost every one, under the name of Indian Pink, and universally used as a vermifuge amongst children; the living vegetable, however, is rarely met with in our gardens.

It seems to have been given up by the faculty for more certain and active medicines. Small doses of the recent plant are said, occasionally, to produce

giddiness, dimness of sight, and other alarming symptoms, whilst larger doses never produce the same effects, on account of its cathartic properties being brought into action.

Several eminent physicians of America, who first introduced the *Spigelia* to notice, have done so under the most favourable impressions of its anthelmintic virtues. One of them, Dr. Gardner, however, observes that he had given it in hundreds of cases, but that he never found its virtues very decided, unless it proved aperient. Dr. Bigelow says the root of this, as of all other perennial plants, is the most active part; and that ten grains may be given to a child four years old.

The small fibrous roots form but an inconsiderable portion of the plant; the entire of which is usually employed in England; and that, always, in a dried state. These circumstances preclude any narcotic effects that may arise from the fresh gathered root. The experience of many medical practitioners has proved that the *Spigelia* is best administered in combination with some more active cathartic medicine, as two or three grains of calomel, or fifteen to twenty grains of rhubarb, for an adult; and less, in proportion, for children.

The best and most popular method of giving it, is in the form of infusion, and combined with senna. Half a dram of each, infused all night in half a tea-cupful of water, kept warm, may be given to a child two or three years old.

It should be planted in a warm, and rather moist, situation, with a portion of peat in the soil; and may occasionally be divided at the root.

Hort. Kew. 2, v. 1, 317.

DIGITALIS LU' TEA.

SMALL YELLOW FOX-GLOVE.

Class.

DIDYNAMIA.

Order.

ANGIOSPERMIA.

Natural Order.

SCROPHULARINÆ.

Native of	Height.	Flowers in	Duration.	Cultivated
S. Europe.	3 feet.	July, Aug.	Perennial.	in 1629.

No. 94.

Digitalis from digitale, the finger of a glove. The name appears first to have been given by the German writer Fuchs, or Fuchsius. Hence the plant was called Digitalis Fuchsii. Sir J. E. Smith suggests that our English appellation, Fox-glove, may have risen from this circumstance.

Most of the species of Fox-glove are extremely showy and attractive plants. The very beautiful spikes of flowers, that are produced by the Digitalis purpurea, or common Fox-glove, and which we so frequently see as a beautiful ornament of waste and high ground, cannot have escaped the notice of the most apathetical observer.

Both the purple and white variety are now frequently met with in gardens and shrubberies, and but few plants exhibit more beauty and gaiety.

The digitalis we may notice, as well for its deleterious and medical qualities, as for its beauty. In the hands of the unskilful it forms an extremely dangerous medicine; whilst Dr. Withering and others have shown, that when administered with skill and caution, it is, perhaps, one of the most valuable vegetables that we possess.

Digitalis is stated as possessing properties which are combined in no other substance. It is a direct sedative, diminishing, most powerfully, the actions of the system, without occasioning previous excitement. Administered with caution, a pulse of 70 beats or more in a minute, will frequently be reduced to 40, or even less. But when thrown into the constitution too suddenly, or if the quantity be too great, it induces vertigo, sickness, convulsions, coldness of the body, extreme debility, and death.

Another powerful quality of this plant, is that of an active diuretic. But when employed as such, great care is required, lest its sudden diminution of the vascular action, should induce so great a degree of debility, as to prove fatal to patients of a weakly or diseased constitution.

Notwithstanding the usual influence of this plant on the human frame, still its powers are not certain; and constitutions have been met with whereon it had no manner of effect, even in excessive doses. In a few cases its effects have not been evinced till its use has been continued for some time; when, at length, its powers have burst forth with the greatest vehemence, so as to endanger the lives of those to whom it was administered.

We have been thus particular, in relating the effects of this plant; and advise that it never be administered but under the direction and superintendence of a medical man. That the unskilful may be informed how small a portion of this deleterious plant is sufficient to act injuriously on the system, we shall briefly state, that of the dried leaves, one to three grains twice a day is a full dose.

Hort. Kew. 2, v. 4, 29.

CENTAUREA SUAVEOLENS.

YELLOW SULTAN.

Class.
SYNGENESIA.

Order.
POLYGAMIA FRUSTRANEA.

Natural Order.
CYNAROCEPHALEÆ.

Native of Levant.	Height. 1 foot.	Flowers in July, Sep.	Duration. Annual.	Cultivated in 1683.
----------------------	--------------------	--------------------------	----------------------	------------------------

No. 95.

Centaurea is derived from a Greek word of similar construction, signifying a centaur. This appellation was given after Chiron the centaur, who is said to have used a species of the Centaurea to cure himself of a wound, occasioned by the falling of one of the arrows of Hercules on his foot. Suaveolens, from the Latin suavis, sweet.

This most elegant and attractive flower may, certainly, be ranked amongst the prettiest of Flora's gifts, bestowed in the form of an annual. It occupies but little room, therefore should be planted near to the hedge of the flower compartment. It is not so hardy as many other annuals, but still requires no great care, provided it be not sown too early in the spring, nor planted in a cold moist situation.

It is sometimes raised on a hot-bed, and transplanted, but when so propagated, it should have a quantity of soil taken up with the roots, and be carefully watered and shaded afterwards.

It does not freely produce seed, unless the head of one flower be shaken over another, by which the pollen is scattered, and the parts of fructification will thereby become fertilized.

Hort. Kew. 2, v. 5, 144.

PRIMULA FARINOSA.

BIRD'S-EYE PRIMROSE.

Class.
PENTANDRIA.

Order.
MONOGYNIA.

Natural Order.
PRIMULACEÆ.

Native of Britain.	Height. 4 inches.	Flowers in June, July.	Duration. Perennial.	Inhabits Wet Mead.
-----------------------	----------------------	---------------------------	-------------------------	-----------------------

No. 96.

The derivation of Primula, from primus, first, has been noticed under No. 1 and 60. Farinosa, from the Latin farina, flour; a beautifully white powdering of which this primula bears on the scape or flower stalk, and on the young leaves, unless it be exposed to rains, and thereby deprived of this now unfashionable appendage.

It is one amongst that beautiful tribe of alpine plants with which every one is pleased, and which every one may possess. It is found in the north of England, in various other parts of Europe, and even in Siberia. In autumn its leaves fade, and the plant is comprised in what may be termed a radical bud, on the surface of the soil. Thus, inhabiting its winter quarters, it remains in security, till the genial warmth of spring expands its mealy foliage, and in due time occasions it again to present us with a lively specimen of beauty in miniature.

The Primula farinosa we find succeed very well, when planted in a small pot of light loam and peat. Severe frosts do it no injury, but it should have a temporary shelter against excessive moisture, during the winter.

Hort. Kew. 2, v. 1, 308.

13 vols
7712



Maund, Benjamin,

The botanic garden Consisting of highly finished representations of hardy
ornamental flowering plants, cultivated in Great Britain ; with their
names, classes, orders, history, qualities, culture and physiological
observations

Bd.: 1. [Nebst] Index 1 - 6. - 1825. - 106 Bl. : 24 Ill.

London 1825

Phyt. 926 I-1

urn:nbn:de:bvb:12-bsb10304263-1